## NO-GAP SECOND-ORDER CONDITIONS VIA A DIRECTIONAL CURVATURE FUNCTIONAL\*

CONSTANTIN CHRISTOF<sup>†</sup> AND GERD WACHSMUTH<sup>‡</sup>

**Abstract.** This paper is concerned with necessary and sufficient second-order conditions for finite-dimensional and infinite-dimensional constrained optimization problems. Using a suitably defined directional curvature functional for the admissible set, we derive no-gap second-order optimality conditions in an abstract functional analytic setting. Our theory not only covers those cases where the classical assumptions of polyhedricity or second-order regularity are satisfied but also allows to study problems in the absence of these requirements. As a tangible example, we consider no-gap second-order conditions for bang-bang optimal control problems.

**Key words.** second-order condition, bang-bang control, polyhedricity, second-order regularity, no-gap optimality condition

AMS subject classifications. 49J53, 49K20, 49K27, 49K30

1. Introduction. The aim of this paper is to develop a theoretical framework for necessary and sufficient second-order optimality conditions (henceforth referred to as SNC and SSC, respectively). The main feature of our approach is that we use a suitably defined "directional" curvature functional to take into account the influence of the admissible set. Our definition of curvature allows us to derive no-gap second-order conditions that provide more flexibility than classical results. In particular, our approach makes it possible to exploit additional information about the gradient of the objective functional. Such information is, e.g., often available in the optimal control of partial differential equations (PDEs) where the gradient of the objective is typically characterized by an adjoint equation and, as a consequence, enjoys additional regularity properties.

Let us briefly clarify what we understand by "no-gap second-order conditions". For simplicity, we focus on the minimization of a smooth function f over  $\mathbb{R}^d$ . In this case, it is well known that local optimality of  $\bar{x}$  implies  $\nabla f(\bar{x}) = 0$  and  $h^\top \nabla^2 f(\bar{x}) \, h \geq 0$  for all  $h \in \mathbb{R}^d$ . On the other hand,  $\nabla f(\bar{x}) = 0$  and  $h^\top \nabla^2 f(\bar{x}) \, h > 0$  for all  $h \in \mathbb{R}^d \setminus \{0\}$  is equivalent to  $\bar{x}$  being a local minimizer satisfying a quadratic growth condition. Hence, the only difference between the necessary and the sufficient condition is a non-strict vs. a strict inequality in the second-order condition. Moreover, this change is as small as possible. Such a pair of optimality conditions is denoted as "no-gap second-order conditions".

The analysis found in this paper originated from the idea to extend the results of [11]. In this paper, the authors derived an SSC for a class of bang-bang optimal control problems that does not fit into the classical setting of polyhedricity and second-order regularity, cf. [4]. Our results turned out to be of relevance for other problems as well and ultimately gave rise to the abstract framework of Section 4 that not only covers large parts of the classical SNC and SSC theory but also allows to study situations where the admissible set exhibits a singular or degenerate curvature behavior, cf. Section 6. We hope that with the subsequent analysis we can, on the one hand, offer

<sup>\*</sup>THIS RESEARCH WAS SUPPORTED BY THE GERMAN RESEARCH FOUNDATION UNDER GRANT NUMBERS ME 3281/7-1 AND WA 3636/4-1 WITHIN THE PRIORITY PROGRAM "NON-SMOOTH AND COMPLEMENTARITY-BASED DISTRIBUTED PARAMETER SYSTEMS: SIMULATION AND HIERARCHICAL OPTIMIZATION" (SPP 1962).

<sup>†</sup>TECHNISCHE UNIVERSITÄT MÜNCHEN, FACULTY OF MATHEMATICS, 85748 GARCHING, GERMANY

<sup>&</sup>lt;sup>‡</sup>BRANDENBURGISCHE TECHNISCHE UNIVERSITÄT COTTBUS-SENFTENBERG, INSTITUTE OF MATHEMATICS, CHAIR OF OPTIMAL CONTROL, 03046 COTTBUS, GERMANY, HTTPS://WWW.B-TU.DE/FG-OPTIMALE-STEUERUNG, GERD.WACHSMUTH@B-TU.DE

an alternative view on well-known SNC and SSC results and, on the other hand, also provide some new ideas for the study of problems that do not satisfy the classical assumptions of polyhedricity and second-order regularity.

Let us give some references: There are several contributions addressing second-order optimality conditions for optimization problems posed in infinite-dimensional spaces. We mention exemplarily [18, 3, 5, 4, 8, 9]. No-gap optimality conditions in the infinite-dimensional case can be found, e.g., in [5, Theorems 2.7, 2.10], [8, Theorems 2.2, 2.3], and [23, Theorem 5.7]. Note that the latter results all rely on the concept of polyhedricity (in contrast to our Theorem 4.5) and require that the Hessian of the Lagrangian is a Legendre form. In the finite-dimensional case, one can further employ the notion of second-order regularity to derive no-gap optimality conditions, see, e.g., [3]. Note that, in our approach, the Legendre form condition is substituted by a more general non-degeneracy condition, see (NDC) in Theorem 4.4 and the discussion in Subsection 5.1.

Before we begin with our analysis, we give a short overview over the contents and the structure of this paper:

In Section 2, we clarify the notation, make our assumptions precise and recall several definitions that are needed for our investigation.

In Section 3, we define the directional curvature functional that is at the heart of our SNC and SSC analysis. Here, we also discuss basic properties of the curvature functional as, e.g., positivity and lower semicontinuity, that are used in the remainder of the paper.

Section 4 addresses SNC and SSC for constrained optimization problems on an abstract functional analytic level. The main results of this section (and of the paper as a whole) are Theorems 4.3 to 4.5. These theorems illustrate the advantages of working with the directional curvature functional and demonstrate that our approach allows for a very short and elegant derivation of the second-order theory.

In Section 5, we demonstrate that the framework of Section 4 indeed covers the classical SNC and SSC theory for optimization problems with polyhedric or second-order regular sets. Here, we further comment on how the assumptions (NDC) and (MRC) appearing in our second-order conditions can be verified in practice and interpreted in the context of generalized Legendre forms and Tikhonov regularization. Section 5 also includes two tangible examples that demonstrate the usefulness of the theorems in Section 4.

Section 6 is devoted to problems that are covered by our analysis but do not fall under the scope of the classical SNC- and SSC-framework. The first example that we consider in this context is a finite-dimensional optimization problem whose admissible set exhibits a singular curvature behavior. In Subsection 6.2, we then study no-gap second-order conditions for bang-bang optimal control problems in the measure space  $\mathcal{M}(\Omega)$ . Here, we prove a novel SNC for bang-bang controls and further sharpen the SSC in [11] to close the gap between the two conditions, see Theorems 6.4 and 6.12, and the comparison in Example 6.14.

Section 7 summarizes our findings and gives some pointers to further research.

**2.** Notation, Preliminaries and Basic Concepts. Throughout this paper, we always consider the following situation.

Assumption 2.1 (Standing Assumptions and Notation).

- (i) X is the (topological) dual of a separable Banach space Y,
- (ii)  $\iota: Y \to X^*$  denotes the canonical embedding of Y into the dual  $X^*$  of X,
- (iii) the admissible set C is a closed, non-empty subset of X.

Note that, under the above assumptions, X is necessarily a Banach space with a weak- $\star$  sequentially compact unit ball, cf. the Banach-Alaoglu theorem. We remark that the overwhelming majority of our results also holds when the space Y is assumed to be reflexive instead of separable. We restrict our analysis to the above setting since it allows to study more interesting practical examples, see Subsection 6.2. For our analysis, we need the following classical concepts (cf. [4, Section 2.2.4]).

DEFINITION 2.2. Let  $x \in C$  be given. We define the radial cone, the strong outer tangent cone and the weak- $\star$  outer tangent cone to C at x, respectively, by

$$\mathcal{R}_{C}(x) := \left\{ h \in X \mid \exists T > 0 \ \forall t \in [0, T], x + th \in C \right\},$$

$$\mathcal{T}_{C}(x) := \left\{ h \in X \mid \exists t_{k} \searrow 0, \exists x_{k} \in C \ such \ that \ \frac{x_{k} - x}{t_{k}} \to h \right\},$$

$$\mathcal{T}_{C}^{\star}(x) := \left\{ h \in X \mid \exists t_{k} \searrow 0, \exists x_{k} \in C \ such \ that \ \frac{x_{k} - x}{t_{k}} \stackrel{\star}{\rightharpoonup} h \right\}.$$

Moreover, we define the weak- $\star$  (outer) normal cone to x by

$$\mathcal{N}_C^{\star}(x) := \{ x^{\star} \in \iota(Y) \mid \forall h \in \mathcal{T}_C^{\star}(x) : \langle x^{\star}, h \rangle \le 0 \}.$$

Finally, given a  $\varphi \in -\mathcal{N}_C^{\star}(x)$ , we define the weak- $\star$  critical cone by

$$\mathcal{K}_C^{\star}(x,\varphi) := \mathcal{T}_C^{\star}(x) \cap \varphi^{\perp}.$$

We emphasize that  $\mathcal{N}_{\mathcal{C}}^{\star}(x)$  is defined to be a subset of  $\iota(Y)$  and may thus be identified with a subset of the predual space Y. Note that all "cones" in the above are indeed cones in the mathematical sense, i.e.,  $h \in \mathcal{R}_{\mathcal{C}}(x)$  implies  $\alpha h \in \mathcal{R}_{\mathcal{C}}(x)$  for all  $\alpha \geq 0$  etc. We point out that  $\mathcal{R}_{\mathcal{C}}(x) \subset \mathcal{T}_{\mathcal{C}}(x) \subset \mathcal{T}_{\mathcal{C}}^{\star}(x)$ . If  $\mathcal{C}$  is convex, then [4, Proposition 2.55]

$$\mathcal{R}_C(x) = \mathbb{R}^+(C - x), \quad \mathcal{T}_C(x) = \operatorname{cl}(\mathcal{R}_C(x)) \quad \forall x \in C.$$

If, in addition, X is reflexive, then we have  $\mathcal{T}_C^{\star}(x) = \mathcal{T}_C(x)$  by Mazur's lemma. We remark that  $\mathcal{T}_C^{\star}(x)$  is in general not closed since the weak- $\star$  topology is not sequential on infinite-dimensional spaces, cf. [19, Example 5.9].

**3.** The Directional Curvature Functional. The basic idea of our SNC and SSC approach is to not discuss the curvature properties of the admissible set C separately, i.e., independently of the optimization problem at hand, but to develop a second-order analysis that takes into account the gradient of the objective. To accomplish the latter, we introduce the directional curvature functional.

Definition 3.1. Let  $x \in C$  and  $\varphi \in -\mathcal{N}_C^{\star}(x)$  be given. The weak- $\star$  directional curvature functional  $Q_C^{x,\varphi} \colon \mathcal{K}_C^{\star}(x,\varphi) \to [-\infty,\infty]$  associated with the triple  $(x,\varphi,C)$  is defined by

$$(1) \quad Q_C^{x,\varphi}(h) := \inf \left\{ \liminf_{k \to \infty} \left\langle \varphi, r_k \right\rangle \, \middle| \, \begin{cases} \{r_k\} \subset X, \{t_k\} \subset \mathbb{R}^+ : t_k \searrow 0, t_k \, r_k \stackrel{\star}{\rightharpoonup} 0, \\ x + t_k \, h + \frac{1}{2} \, t_k^2 \, r_k \in C \end{cases} \right\}.$$

In case that X is finite-dimensional, the curvature functional  $Q_C^{x,\varphi}$  coincides with a generalized derivative of second order of the indicator function  $\delta_C: X \to \{0, \infty\}$ ,

which is termed "second subderivative" in [21, Definition 13.3]. A similar concept in infinite dimensions is called "second-order epiderivative" in [14, Section 1]. Therein, it is required that the second-order difference quotients associated with the indicator function of the set C, the point x, and the element  $\varphi \in -\mathcal{N}_{\mathcal{L}}^{\star}(x)$  Mosco epi-converge, whereas our definition just uses the weak-\* limes superior (in the sense of Kuratowski) of the epigraphs of the difference quotients, see also Remark 3.5 below. The functional  $Q_C^{x,\varphi}(\cdot)$  may thus be identified with a second-order weak-\* lower subderivative.

Definition 3.1 can be motivated as follows: Consider an optimization problem of the form min  $J(x) := \langle \varphi, x \rangle$  s.t.  $x \in C$ , where  $\varphi$  satisfies  $\varphi \in -\mathcal{N}_C^{\star}(\bar{x})$  for some  $\bar{x} \in C$ , i.e.,  $\langle \varphi, h \rangle \geq 0$  for all  $h \in \mathcal{T}_C^{\star}(\bar{x})$ . Then  $\bar{x}$  is a critical point and we have to study perturbations into the critical directions  $h \in \mathcal{K}_C^{\star}(\bar{x},\varphi)$  to decide whether  $\bar{x}$  is a local minimizer or not. If we fix a critical direction  $h \in \mathcal{K}_C^{\star}(\bar{x}, \varphi)$ , then the definition of  $\mathcal{T}_C^{\star}(\bar{x})$  yields the existence of sequences  $\{r_k\} \subset X$ ,  $\{t_k\} \subset \mathbb{R}^+$  with  $t_k \searrow 0$ ,  $t_k r_k \stackrel{\star}{\searrow} 0$  and  $x_k := \bar{x} + t_k h + \frac{1}{2} t_k^2 r_k \in C$ , and we may calculate that  $J(x_k) - J(\bar{x}) = \frac{1}{2} t_k^2 \langle \varphi, r_k \rangle$ . This identity suggests that the limiting behavior of the dual pairing  $\langle \varphi, r_k \rangle$  between the gradient  $\varphi$  of the objective J and the second-order correction  $r_k$  is a decisive factor in the study of the optimality of the critical point  $\bar{x}$ . Since the directional curvature functional  $Q_C^{x,\varphi}(\cdot)$  allows to estimate the limes inferior of precisely that quantity for all possible sequences  $\{r_k\}$  and  $\{t_k\}$ , it is only natural to consider it an adequate tool for the derivation of optimality conditions. Note that, instead of looking at the accumulation points of the corrections  $r_k$ , which is the idea of secondorder tangent sets, cf. Definition 5.2, we only study accumulation points of the scalar sequences  $\langle \varphi, r_k \rangle$  when working with the functional  $Q_C^{\bar{x}, \varphi}(\cdot)$ . Thus, it is possible to obtain information even when the second-order corrections  $r_k$  diverge or cannot be analyzed properly. Before turning our attention to SNC and SSC, in what follows, we first state some preliminary results on the properties of the directional curvature functional that are needed for our investigation.

Lemma 3.2. Let  $x \in C$  be given. Then the following assertions hold.

- (i) For all  $\varphi \in -\mathcal{N}_C^{\star}(x)$ ,  $h \in \mathcal{K}_C^{\star}(x,\varphi)$ ,  $\alpha > 0$ , we have  $Q_C^{x,\varphi}(\alpha h) = \alpha^2 Q_C^{x,\varphi}(h)$ . (ii) If C is convex, then  $Q_C^{x,\varphi}(h) \geq 0$  for all  $\varphi \in -\mathcal{N}_C^{\star}(x)$  and  $h \in \mathcal{K}_C^{\star}(x,\varphi)$ .

Proof. Assertion (i) can be checked by a simple scaling argument. To prove (ii), suppose that C is convex, let  $h \in \mathcal{K}_C^{\star}(x,\varphi)$  and  $\varphi \in -\mathcal{N}_C^{\star}(x)$  be arbitrary but fixed, and let  $\{r_k\}$ ,  $\{t_k\}$  be sequences as in the definition of  $Q_C^{x,\varphi}(h)$ . Then it holds  $\frac{2}{t_k}h + r_k \in \mathcal{R}_C(x) \subset \mathcal{T}_C^*(x)$  due to the convexity of C and, consequently,  $\langle \varphi, r_k \rangle = \langle \varphi, \frac{2}{t_k}h + r_k \rangle \geq 0$ . Taking the limes inferior for  $k \to \infty$  and the infimum over all  $\{r_k\}$ ,  $\{t_k\}$  now yields the claim. 

In addition to Lemma 3.2, we have the following weak-\* lower semicontinuity result.

LEMMA 3.3. Let  $x \in C$  and  $\varphi \in -\mathcal{N}_C^{\star}(x)$  be given. Let  $\{h_n\} \subset \mathcal{K}_C^{\star}(x,\varphi)$  be a sequence such that  $h_n \stackrel{\star}{\rightharpoonup} h$  holds for some  $h \in X$  and such that there exist sequences  $\{r_{n,k}\}\subset X \text{ and } \{t_{n,k}\}\subset \mathbb{R}^+ \text{ and a constant } M>0 \text{ with }$ 

$$t_{n,k} \searrow 0, \quad t_{n,k} r_{n,k} \stackrel{\star}{\rightharpoonup} 0, \quad \langle \varphi, r_{n,k} \rangle \to Q_C^{x,\varphi}(h_n) \quad \text{for all $n$ as $k \to \infty$ and}$$
$$\|t_{n,k} r_{n,k}\|_X \leq M, \quad x + t_{n,k} h_n + \frac{1}{2} t_{n,k}^2 r_{n,k} \in C \quad \text{for all $n,k$}.$$

Then, h is an element of the critical cone  $\mathcal{K}_{C}^{\star}(x,\varphi)$  and it holds

(2) 
$$Q_C^{x,\varphi}(h) \le \liminf_{n \to \infty} Q_C^{x,\varphi}(h_n).$$

*Proof.* Consider a countable dense subset  $\{y_i\}$  of Y and choose a sequence  $k_n$  such that

(3) 
$$t_{n,k_n} \leq \frac{1}{n}$$
,  $\langle \varphi, r_{n,k_n} \rangle \leq Q_C^{x,\varphi}(h_n) + \frac{1}{n}$  and  $|\langle y_i, t_{n,k_n}, r_{n,k_n} \rangle| \leq \frac{1}{n}$   $\forall i \leq n$ 

holds for all  $n \in \mathbb{N}$ . Then, our assumptions on  $r_{n,k}$  imply  $||t_{n,k_n} r_{n,k_n}||_X \leq M$ , and we may deduce from (3) that the sequences  $t_n := t_{n,k_n}$  and  $r_n := r_{n,k_n} + \frac{2}{t_n} (h_n - h)$  satisfy

$$t_n \searrow 0, \quad t_n r_n \stackrel{\star}{\rightharpoonup} 0, \quad x + t_n \, h + \frac{1}{2} \, t_n^2 \, r_n \in C \quad \text{and} \quad \liminf_{n \to \infty} \langle \varphi, r_n \rangle \leq \liminf_{n \to \infty} Q_C^{x, \varphi}(h_n).$$

The above yields  $h \in \mathcal{T}_C^{\star}(x)$  and, since we trivially have  $h \in \varphi^{\perp}$ ,  $h \in \mathcal{K}_C^{\star}(x,\varphi)$ . Moreover, we obtain (2) from the definition of  $Q_C^{x,\varphi}(h)$ . This proves the claim.

We point out that Lemma 3.3 is in particular applicable if for every  $h \in \mathcal{K}_C^{\star}(x,\varphi)$  we can find sequences  $\{r_k\} \subset X$  and  $\{t_k\} \subset \mathbb{R}^+$  that realize the infimum in (1) with strong convergence  $t_k r_k \to 0$ . Since sets C with the latter property prove to be useful also in different contexts, we introduce the following concept.

DEFINITION 3.4 (Mosco Regularity Condition (MRC)). We say that C is Mosco regular in  $(x, \varphi) \in C \times -\mathcal{N}_C^{\star}(x)$  if

$$(MRC) \forall h \in \mathcal{K}_C^{\star}(x,\varphi) \; \exists \{r_k\} \subset X, \{t_k\} \subset \mathbb{R}^+ : \\ t_k \searrow 0, \; t_k \, r_k \to 0, \; x + t_k \, h + \frac{1}{2} \, t_k^2 \, r_k \in C, \; Q_C^{x,\varphi}(h) = \lim_{k \to \infty} \left\langle \varphi, r_k \right\rangle.$$

REMARK 3.5. It is easy to see that (MRC) holds in  $(x, \varphi) \in C \times -\mathcal{N}_C^{\star}(x)$  if and only if

$$Q_C^{x,\varphi}(h) = \inf \left\{ \liminf_{k \to \infty} \left\langle \varphi, r_k \right\rangle \, \middle| \, \begin{cases} \{r_k\} \subset X, \{t_k\} \subset \mathbb{R}^+ : t_k \searrow 0, t_k \, r_k \to 0, \\ x + t_k \, h + \frac{1}{2} \, t_k^2 \, r_k \in C \end{cases} \right\}$$

for all  $h \in \mathcal{K}_C^{\star}(x,\varphi)$ , i.e., if and only if the functional  $Q_C^{x,\varphi}(\cdot)$  remains unchanged when we replace the weak- $\star$  convergence of  $t_k r_k$  with strong convergence. In the context of Kuratowski limits, the latter means that the weak- $\star$  limes superior and the strong limes superior of the epigraphs of the second-order difference quotients associated with the indicator function of the set C, the point x, and the element  $\varphi \in -\mathcal{N}_C^{\star}(x)$  coincide. We again refer to [14] for details.

We will see in the following section that the condition (MRC) is also of significance for the study of second-order optimality conditions as it allows to weaken the regularity assumptions on the objective needed for the derivation of SNC. Note that (MRC) is always satisfied when X is finite-dimensional. Further conditions ensuring (MRC) can be found in Lemmas 5.3 and 5.7.

4. Necessary and Sufficient Second-Order Conditions. Having introduced the directional curvature functional  $Q_C^{x,\varphi}(\cdot)$ , we now turn our attention to SNC and SSC for minimization problems of the form

(P) Minimize 
$$J(x)$$
, such that  $x \in C$ .

In the remainder of this paper, when discussing optimality conditions for a problem of the type (P), we always require that (in addition to our standing Assumption 2.1) the following assumption holds.

Assumption 4.1.

- (i)  $\bar{x}$  is a fixed element of the set C (the minimizer/candidate for a minimizer),
- (ii)  $J: C \to \mathbb{R}$  is a function such that there exist a  $J'(\bar{x}) \in \iota(Y)$  and a bounded bilinear  $J''(\bar{x}): X \times X \to \mathbb{R}$  with

(4) 
$$\lim_{k \to \infty} \frac{J(\bar{x} + t_k h_k) - J(\bar{x}) - t_k J'(\bar{x}) h_k - \frac{1}{2} t_k^2 J''(\bar{x}) h_k^2}{t_k^2} = 0$$

for all  $\{h_k\} \subset X$ ,  $\{t_k\} \subset \mathbb{R}^+$  satisfying  $t_k \searrow 0$ ,  $h_k \stackrel{\star}{\rightharpoonup} h \in X$  and  $\bar{x} + t_k h_k \in C$ .

Note that we use the abbreviations  $J'(\bar{x}) h := \langle J'(\bar{x}), h \rangle$  and  $J''(\bar{x}) h^2 := J''(\bar{x})(h, h)$  for all  $h \in X$  in (4), and that (4) is automatically satisfied if J admits a second-order Taylor expansion of the form

(5) 
$$J(\bar{x}+h) - J(\bar{x}) - J'(\bar{x})h - \frac{1}{2}J''(\bar{x})h^2 = o(\|h\|_X^2) \quad \text{as } \|h\|_X \to 0.$$

We begin our investigation by stating necessary conditions of first order.

Theorem 4.2 (First-Order Necessary Condition). Suppose that  $\bar{x}$  is a local minimizer of (P), i.e., assume that there is an  $\varepsilon > 0$  with

$$J(x) \ge J(\bar{x}) \qquad \forall x \in C \cap B_{\varepsilon}^X(\bar{x}),$$

where  $B_{\varepsilon}^{X}(\bar{x})$  denotes the closed ball of radius  $\varepsilon$  around  $\bar{x}$ . Then,  $J'(\bar{x}) \in -\mathcal{N}_{C}^{\star}(\bar{x})$ .

*Proof.* Let  $h \in \mathcal{T}_C^{\star}(\bar{x})$  be given. By definition, there are sequences  $\{x_k\} \subset C$ ,  $\{t_k\} \subset \mathbb{R}^+$  with  $t_k \searrow 0$  and  $(x_k - \bar{x})/t_k \stackrel{\star}{\rightharpoonup} h$ . We set  $h_k := (x_k - \bar{x})/t_k$ . Then for large enough k, we have (due to (4) and the boundedness of weakly- $\star$  convergent sequences)

$$0 \le \frac{J(x_k) - J(\bar{x})}{t_k} = \frac{J(\bar{x} + t_k h_k) - J(\bar{x})}{t_k} = J'(\bar{x}) h_k + \mathcal{O}(t_k) \to J'(\bar{x}) h.$$

The above and our assumption  $J'(\bar{x}) \in \iota(Y)$  yield  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$  as claimed. Using Theorem 4.2, we can provide second-order optimality conditions for (P).

Theorem 4.3 (SNC Involving the Directional Curvature Functional). Suppose that  $\bar{x}$  is a local minimizer of (P) such that

(6) 
$$J(x) \ge J(\bar{x}) + \frac{c}{2} \|x - \bar{x}\|_X^2 \qquad \forall x \in C \cap B_{\varepsilon}^X(\bar{x})$$

holds for some  $c \geq 0$  and some  $\varepsilon > 0$ . Assume further that one of the following conditions is satisfied.

- (i) The map  $h \mapsto J''(\bar{x}) h^2$  is weak-\* upper semicontinuous.
- (ii) The admissible set C satisfies (MRC) in  $(\bar{x}, J'(\bar{x})) \in C \times -\mathcal{N}_C^{\star}(\bar{x})$ . Then

(7) 
$$Q_C^{\bar{x},J'(\bar{x})}(h) + J''(\bar{x}) h^2 \ge c \|h\|_X^2 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{x},J'(\bar{x})).$$

*Proof.* We first consider the case with (i): From the definition of  $Q_C^{\bar{x},J'(\bar{x})}(\cdot)$ , it follows that for every  $h \in \mathcal{K}_C^{\star}(\bar{x},J'(\bar{x}))$  and every  $\delta > 0$  we can find sequences  $\{r_k\} \subset X$  and  $\{t_k\} \subset \mathbb{R}^+$  such that  $t_k \searrow 0$ ,  $t_k r_k \stackrel{\star}{\rightharpoonup} 0$ ,  $x_k := \bar{x} + t_k h + \frac{1}{2} t_k^2 r_k \in C$  and

(8) 
$$\lim_{k \to \infty} \langle J'(\bar{x}), r_k \rangle \le Q_C^{\bar{x}, J'(\bar{x})}(h) + \delta.$$

Since  $x_k \to \bar{x}$  strongly in X, (6) entails  $J(x_k) \geq J(\bar{x}) + \frac{c}{2} \|x_k - \bar{x}\|_X^2$  for large enough k. Hence, we may use (4),  $J'(\bar{x})h = 0$  and the weak- $\star$  lower semicontinuity of the norm  $\|\cdot\|_X$  to obtain

$$0 = \lim_{k \to \infty} \frac{J(x_k) - J(\bar{x}) - t_k J'(\bar{x}) \left(h + \frac{1}{2} t_k r_k\right) - \frac{1}{2} t_k^2 J''(\bar{x}) \left(h + \frac{1}{2} t_k r_k\right)^2}{t_k^2}$$

$$(9) \qquad \geq \limsup_{k \to \infty} \frac{\frac{c}{2} t_k^2 \|h + \frac{1}{2} t_k r_k\|_X^2 - t_k J'(\bar{x}) \left(\frac{1}{2} t_k r_k\right) - \frac{1}{2} t_k^2 J''(\bar{x}) \left(h + \frac{1}{2} t_k r_k\right)^2}{t_k^2}$$

$$\geq \frac{c}{2} \|h\|_X^2 - \frac{1}{2} \liminf_{k \to \infty} \langle J'(\bar{x}), r_k \rangle - \frac{1}{2} \limsup_{k \to \infty} \left(J''(\bar{x}) \left(h + \frac{1}{2} t_k r_k\right)^2\right).$$

From (8), (9) and the weak-\* upper semicontinuity of  $h \mapsto J''(\bar{x}) h^2$ , it follows

$$c||h||_X^2 \le Q_C^{\bar{x},J'(\bar{x})}(h) + \delta + J''(\bar{x})h^2.$$

Passing to the limit  $\delta \searrow 0$  in this inequality yields the claim in the first case.

It remains to prove (7) under assumption (ii). To this end, we note that, if (MRC) holds in  $(\bar{x}, J'(\bar{x})) \in C \times -\mathcal{N}_C^{\star}(\bar{x})$ , then for every  $h \in \mathcal{K}_C^{\star}(\bar{x}, J'(\bar{x}))$  we can find  $\{r_k\} \subset X, \{t_k\} \subset \mathbb{R}^+$  such that  $t_k \searrow 0$ ,  $t_k r_k \to 0$ ,  $x_k := \bar{x} + t_k h + \frac{1}{2} t_k^2 r_k \in C$  and

$$\lim_{k \to \infty} \langle J'(\bar{x}), r_k \rangle = Q_C^{\bar{x}, J'(\bar{x})}(h).$$

Now, the second-order condition (7) follows analogously to case (i).

THEOREM 4.4 (SSC Involving the Directional Curvature Functional). Assume that the map  $h \mapsto J''(\bar{x}) h^2$  is weak-\* lower semicontinuous, that  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$  and that

(10) 
$$Q_C^{\bar{x},J'(\bar{x})}(h) + J''(\bar{x}) h^2 > 0 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{x},J'(\bar{x})) \setminus \{0\}.$$

Suppose further that

(NDC) for all 
$$\{h_k\} \subset X$$
,  $\{t_k\} \subset \mathbb{R}^+$  with  $\bar{x} + t_k h_k \in C$ ,  $h_k \stackrel{\star}{\rightharpoonup} 0$ ,  $t_k \searrow 0$  and  $||h_k||_X = 1$ , it is true that 
$$\lim_{k \to \infty} \inf \left( \langle J'(\bar{x}), h_k/t_k \rangle + \frac{1}{2} J''(\bar{x}) h_k^2 \right) > 0.$$

Then  $\bar{x}$  satisfies the growth condition (6) with some constants c > 0 and  $\varepsilon > 0$ .

*Proof.* We argue by contradiction. Assume that there are no c > 0,  $\varepsilon > 0$  such that (6) holds. Then there are sequences  $\{x_k\} \subset C$  and  $\{c_k\} \subset \mathbb{R}^+$  such that

$$c_k \searrow 0$$
,  $\|x_k - \bar{x}\|_X \to 0$ , and  $J(x_k) < J(\bar{x}) + \frac{c_k}{2} \|x_k - \bar{x}\|_X^2$ .

Define  $t_k := \|x_k - \bar{x}\|_X$  and  $h_k := (x_k - \bar{x})/t_k$ . Then  $\|h_k\|_X = 1$  for all k and we may

extract a subsequence (not relabeled) such that  $h_k \stackrel{\star}{\rightharpoonup} h \in \mathcal{T}_C^{\star}(\bar{x})$ . From (4), it follows

$$0 = \lim_{k \to \infty} \frac{J(\bar{x} + t_k h_k) - J(\bar{x}) - t_k J'(\bar{x}) h_k - \frac{1}{2} t_k^2 J''(\bar{x}) h_k^2}{t_k^2}$$

$$\leq \liminf_{k \to \infty} \frac{\frac{c_k}{2} \|x_k - \bar{x}\|_X^2 - t_k J'(\bar{x}) h_k - \frac{1}{2} t_k^2 J''(\bar{x}) h_k^2}{t_k^2}$$

$$= \liminf_{k \to \infty} \frac{-J'(\bar{x}) h_k - \frac{1}{2} t_k J''(\bar{x}) h_k^2}{t_k}$$

$$\leq -\limsup_{k \to \infty} \frac{J'(\bar{x}) h_k}{t_k} - \liminf_{k \to \infty} \frac{\frac{1}{2} t_k J''(\bar{x}) h_k^2}{t_k}$$

$$\leq -\limsup_{k \to \infty} \frac{J'(\bar{x}) h_k}{t_k} - \frac{1}{2} J''(\bar{x}) h^2.$$

Thus,  $\limsup_{k\to\infty} J'(\bar{x}) h_k/t_k$  is bounded from above and from  $t_k \searrow 0$  we infer  $\limsup_{k\to\infty} J'(\bar{x}) h_k \leq 0$ . Together with  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$  we find  $J'(\bar{x}) h = 0$ . Using  $h \in \mathcal{T}_C^{\star}(\bar{x})$ , this yields  $h \in \mathcal{K}_C^{\star}(\bar{x}, J'(\bar{x}))$ . Setting  $r_k := 2(h_k - h)/t_k$  and using (11),  $J'(\bar{x}) h = 0$  and Definition 3.1, we obtain

$$J''(\bar{x}) h^2 \le -2 \limsup_{k \to \infty} \frac{J'(\bar{x}) h_k}{t_k} = -\limsup_{k \to \infty} J'(\bar{x}) r_k \le -\liminf_{k \to \infty} J'(\bar{x}) r_k$$
  
$$\le -Q_C^{\bar{x}, J'(\bar{x})}(h).$$

From (10), we may now deduce that h is zero. This is a contradiction with (NDC), see the properties of the sequences  $\{h_k\}$ ,  $\{t_k\}$  and (11).

The acronym (NDC) stands for "non-degeneracy condition". Comments on (NDC) are provided in Subsection 5.1. By combining the previous two theorems, we arrive at our main theorem on no-gap second-order conditions.

THEOREM 4.5 (No-Gap Second-Order Optimality Condition). Assume that the map  $h \mapsto J''(\bar{x}) h^2$  is weak-\* lower semicontinuous, that  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$ , that (NDC) holds, and that one of the conditions (i) and (ii) in Theorem 4.3 is satisfied. Then, the condition

$$Q_C^{\bar{x},J'(\bar{x})}(h) + J''(\bar{x}) h^2 > 0 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{x},J'(\bar{x})) \setminus \{0\}$$

is equivalent to the quadratic growth condition (6) with constants c > 0 and  $\varepsilon > 0$ .

Some remarks regarding Theorems 4.3 to 4.5 are in order.

## Remark 4.6.

- (i) Note that Theorem 4.3 yields that there are two ways to obtain the secondorder necessary condition (7) in the situation of Assumption 4.1: We can assume either that the second derivative of J at  $\bar{x}$  has additional (semi)continuity properties (this is case (i)) or that the set C is sufficiently wellbehaved at  $\bar{x}$  (this is case (ii)). The observation that one has the choice to impose additional assumptions either on the appearing sets or the appearing functions can be made frequently when working with the directional curvature functional.
- (ii) We point out that the regularity condition (MRC) is not helpful in the derivation of the SSC in Theorem 4.4. At the heart of the proof of Theorem 4.4 is,

- after all, the contradiction argument which only provides a weak-⋆ convergent subsequence.
- (iii) It is easy to see that the differentiability assumptions on J in Assumption 4.1 can be weakened when one is interested in only one of the results in Theorems 4.3 and 4.4. To derive the necessary condition (7) in case (ii) of Theorem 4.3, for example, it suffices to assume that

$$h \in X, \{r_k\} \subset X, \ \{t_k\} \subset \mathbb{R}^+, t_k \searrow 0, t_k r_k \to 0 \ in \ X$$

$$\Rightarrow \max\left(0, \frac{J(\bar{x} + t_k h + \frac{1}{2}t_k^2 r_k) - J(\bar{x}) - t_k J'(\bar{x})(h + \frac{1}{2}t_k r_k) - \frac{1}{2}t_k^2 J''(\bar{x})h^2}{t_k^2}\right) \to 0$$

- holds for some bounded, linear mapping  $J'(\bar{x}): X \to \mathbb{R}$  and some bilinear form  $J''(\bar{x}): X \times X \to \mathbb{R}$ . We refrain from stating the minimal differentiability properties in each of the Theorems 4.3 to 4.5 since this would just obscure the basic ideas of our analysis.
- (iv) We point out that it is possible to modify the proofs of Theorems 4.3 and 4.4 to obtain second-order conditions in the setting of two-norms discrepancy, i.e., in the situation where the objective J only satisfies a second-order Taylor expansion à la (5) w.r.t. some norm  $\|\cdot\|_Z$  that is stronger than  $\|\cdot\|_X$ . We leave it to the interested reader to work out this easy extension of our analysis in detail.
- (v) In the finite-dimensional setting, results similar to Theorems 4.3 to 4.5 have been obtained in [21, Theorem 13.24].
- 5. How to Apply and Interpret the Results of Section 4 in the Context of the Classical Theory. In this section, we comment on the verification and interpretation of the conditions (NDC) and (MRC) appearing in our second-order conditions, see Subsection 5.1, Lemma 5.3, and Lemma 5.7. Further, we address the computation of the directional curvature functional for polyhedric and second-order regular sets, see Subsection 5.2. Finally, we compare our theorems of Section 4 with classical results. In Subsection 5.3, we use the observations of Subsections 5.1 and 5.2 to state two corollaries of Theorem 4.5 that reproduce (and slightly extend) classical results found, e.g., in [4] and [9]. We conclude this section with two examples that illustrate the usefulness of Theorems 4.3 to 4.5.
- **5.1.** Remarks on the Non-Degeneracy Condition (NDC). The condition (NDC) appearing in our SSC can be interpreted as a generalized Legendre condition, cf. [4, Section 3.3.2] and Lemma 5.1 (ii). In contrast to a classical Legendre condition, our requirement (NDC) is a condition on the interplay between the curvature of Cin  $\bar{x}$  in the direction  $J'(\bar{x})$  and the properties of  $J''(\bar{x})$ . In practice, (NDC) can be ensured, e.g., by assuming ellipticity of the second derivative  $J''(\bar{x})$  or by assuming that the admissible set C has "positive" curvature at  $\bar{x}$  (or some combination of the both). Some sufficient criteria can be found in the following lemma.
  - LEMMA 5.1. Each of the following conditions is sufficient for (NDC).
  - (i) C is a convex subset of X,  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$ , and  $J''(\bar{x})h^2 \geq r > 0$  for all  $h \in X$  with  $||h||_X = 1$ .
  - (ii) C is a convex subset of X,  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$ , and  $J''(\bar{x})$  is a Legendre form in the sense that  $h \mapsto J''(\bar{x}) h^2$  is weak-\* lower semicontinuous and

$$h_k \stackrel{\star}{\rightharpoonup} h \ and \ J''(\bar{x}) h_k^2 \to J''(\bar{x}) h^2 \quad \Rightarrow \quad h_k \to h$$

holds for all sequences  $\{h_k\} \subset X$ .

- (iii) There exist  $c, \varepsilon > 0$  with  $J'(\bar{x})(x \bar{x}) \ge \frac{c}{2} ||x \bar{x}||^2$  for all  $x \in B_{\varepsilon}^X(\bar{x}) \cap C$  and the map  $h \mapsto J''(\bar{x})h^2$  is weak-\* lower semicontinuous.
- (iv) X is finite-dimensional.

*Proof.* Case (iv) is trivial. To prove (NDC) in the cases (i) to (iii), we have to show that for all  $\{t_k\}$ ,  $\{h_k\}$  satisfying  $t_k \searrow 0$ ,  $h_k \stackrel{\star}{\rightharpoonup} 0$  and  $\|h_k\|_X = 1$ ,  $\bar{x} + t_k h_k \in C$  for all k, it holds  $\lim\inf_{k\to\infty} \left(\langle J'(\bar{x}), h_k/t_k \rangle + \frac{1}{2}J''(\bar{x}) h_k^2\right) > 0$ . Therefore, we assume that sequences  $\{t_k\}$ ,  $\{h_k\}$  with the above properties are given.

We first discuss the case that C is convex. For such a C, it holds  $h_k \in \mathcal{R}_C(\bar{x})$  and, as a consequence,  $J'(\bar{x}) h_k \geq 0$ , so it suffices to prove  $\liminf_{k \to \infty} J''(\bar{x}) h_k^2 > 0$  to obtain the claim. The latter is trivially true under assumption (i). Similarly, in case (ii), we have  $\liminf_{k \to \infty} J''(\bar{x}) h_k^2 \geq 0$  by lower semicontinuity. Moreover, if we had  $J''(\bar{x}) h_k^2 \to 0$  (along a subsequence), we would obtain the contradiction  $h_k \to 0$  (along a subsequence). Hence,  $\liminf_{k \to \infty} J''(\bar{x}) h_k^2 > 0$  and (NDC) is proved for (ii).

In case (iii), we have

$$\langle J'(\bar{x}), h_k/t_k \rangle = t_k^{-2} \langle J'(\bar{x}), t_k h_k \rangle \ge \frac{c}{2} t_k^{-2} \|t_k h_k\|^2 = \frac{c}{2}$$

for k large enough. Together with  $\liminf_{k\to\infty}J''(\bar x)\,h_k^2\geq J''(\bar x)\,0^2=0$ , the above yields the desired inequality  $\liminf_{k\to\infty}\left(\langle J'(\bar x),h_k/t_k\rangle+\frac12\,J''(\bar x)\,h_k^2\right)>0$ . This completes the proof.

Note that in case (iii) of Lemma 5.1, the second derivative  $J''(\bar{x})$  is allowed to be negative definite. This effect occurs whenever the curvature of the set C at  $\bar{x}$  is such that it can compensate for negative curvature of the objective J. In Subsection 6.2, we will see an example, where the latter actually happens. We further point out that, under the assumptions of Lemma 5.1 (iii), the directional curvature functional is coercive in the sense that

(12) 
$$Q_C^{\bar{x},J'(\bar{x})}(h) \ge c \|h\|_X^2 \qquad \forall h \in \mathcal{K}_C^{\star}(\bar{x},J'(\bar{x})),$$

where c is the constant in Lemma 5.1 (iii). Indeed, for  $h \in \mathcal{K}_C^{\star}(\bar{x}, J'(\bar{x}))$ ,  $t_k \searrow 0$ ,  $t_k r_k \stackrel{\star}{\rightharpoonup} 0$  and  $x_k := \bar{x} + t_k h + \frac{1}{2} t_k^2 r_k \in C$ , we have

$$\liminf_{k \to \infty} J'(\bar{x}) \, r_k = \liminf_{k \to \infty} J'(\bar{x}) \, \frac{x_k - \bar{x}}{\frac{1}{2} \, t_k^2} \\
\geq \liminf_{k \to \infty} c \, \frac{\|x_k - \bar{x}\|_X^2}{t_k^2} = \liminf_{k \to \infty} c \, \|h + \frac{1}{2} \, t_k \, r_k\|_X^2 \geq c \, \|h\|_X^2.$$

The above gives a lower estimate for the directional curvature functional that is often also interesting for its own sake. We will get back to this topic in Subsection 6.2.

**5.2.** Directional Curvature, Polyhedricity and Second-Order Regularity. In this section, we discuss how the notion of directional curvature is related to the classical concepts of polyhedricity and second-order regularity, and how these properties can be used to calculate the functional  $Q_C^{x,\varphi}(\cdot)$  for a given tuple  $(x,\varphi) \in C \times -\mathcal{N}_C^{\star}(x)$ . Recall the following definitions, cf. [4, Section 3.2.1, Definitions 3.51, 3.85], [20], and [23, Lemma 4.1].

Definition 5.2 (Polyhedricity and Second-Order Regularity).

(i) The set C is said to be polyhedric at  $x \in C$  if C is convex and

$$\mathcal{T}_C(x) \cap \varphi^{\perp} = \operatorname{cl}\left(\mathcal{R}_C(x) \cap \varphi^{\perp}\right) \quad \forall \varphi \in X^{\star}.$$

(ii) The strong (outer) second-order tangent set to a tuple  $(x,h) \in C \times \mathcal{T}_C(x)$  is given by

$$\mathcal{T}^2_C(x,h) := \left\{ r \in X \mid \exists t_n \searrow 0, \operatorname{dist}\left(x + t_n h + \frac{1}{2} t_n^2 r, C\right) = o(t_n^2) \right\}.$$

(iii) The set C is called (outer) second-order regular at  $x \in C$  if for all  $h \in \mathcal{T}_C(x)$  and all  $x_n \in C$  of the form  $x_n := x + t_n h + \frac{1}{2}t_n^2r_n$  with  $t_n \searrow 0$  and  $t_n r_n \to 0$  it holds

$$\lim_{n \to \infty} \operatorname{dist} \left( r_n, \mathcal{T}_C^2(x, h) \right) = 0.$$

Note that sequences  $\{x_n\}$ ,  $\{r_n\}$  as in (iii) exist for all  $h \in \mathcal{T}_C(x)$  and that a set C can only be (outer) second-order regular at  $x \in C$  if  $\mathcal{T}_C^2(x,h) \neq \emptyset$  for all  $h \in \mathcal{T}_C(x)$  (since otherwise  $\operatorname{dist}(\cdot, \mathcal{T}_C^2(x,h)) = +\infty$  by the usual conventions).

First, we show that the boundary of polyhedric sets is not curved.

LEMMA 5.3 (Curvature of Polyhedric Sets). Assume that X is reflexive and that C is polyhedric at  $x \in C$ . Then (MRC) is satisfied in  $(x, \varphi)$  for all  $\varphi \in -\mathcal{N}_C^{\star}(x)$  and

$$Q_C^{x,\varphi}(h) = 0 \qquad \forall h \in \mathcal{K}_C^{\star}(x,\varphi).$$

Proof. From the reflexivity, Mazur's lemma and the convexity and closedness of C, it follows  $\mathcal{T}_C^{\star}(x) = \mathcal{T}_C(x)$ . Consequently,  $\mathcal{K}_C^{\star}(x,\varphi) = \mathcal{T}_C(x) \cap \varphi^{\perp}$  for every  $\varphi \in -\mathcal{N}_C^{\star}(x)$ . For  $h \in \mathcal{R}_C(x) \cap \varphi^{\perp}$  the choice  $t_k = 1/k$ ,  $k \in \mathbb{N}$  sufficiently large, and  $r_k = 0$  shows  $Q_C^{x,\varphi}(h) = 0$ , see also Lemma 3.2 (ii). Now, let  $h \in \mathcal{K}_C^{\star}(x,\varphi) = \mathcal{T}_C(x) \cap \varphi^{\perp}$  be given. Owing to the polyhedricity of C at x, there exists a sequence  $\{h_k\} \subset \mathcal{R}_C(x) \cap \varphi^{\perp}$  with  $h_k \to h$  in X. Now, we can apply Lemma 3.2 (ii) and Lemma 3.3 (with M = 0) to obtain

$$0 \le Q_C^{x,\varphi}(h) \le \liminf_{k \to \infty} Q_C^{x,\varphi}(h_k) = 0.$$

The recovery sequence in (MRC) can be constructed straightforwardly from  $\{h_k\}$ .

Lemma 5.3 shows that, for a polyhedric set C, the directional curvature functional is always identical zero. This is, of course, exactly what one would expect (cf. also with [14, Example 2.10] in this context). The situation is different when C possesses curvature in the sense of second-order regularity – a concept that is promoted and extensively used in [4]. The curvature of second-order regular sets is addressed in the next lemma.

LEMMA 5.4 (Curvature of Second-Order Regular Sets). Assume that C is outer second-order regular at  $x \in C$ , that  $\varphi \in -\mathcal{N}_C^{\star}(x)$ , and that (MRC) is satisfied in  $(x, \varphi)$ . Then

$$Q_C^{x,\varphi}(h) = \inf_{r \in \mathcal{T}_C^2(x,h)} \langle \varphi, r \rangle \qquad \forall h \in \mathcal{K}_C^{\star}(x,\varphi).$$

*Proof.* Let  $h \in \mathcal{K}_C^{\star}(x, \varphi)$  be given. Then (MRC) yields that we can find sequences  $\{r_k\} \subset X$ ,  $\{t_k\} \subset \mathbb{R}^+$  such that  $t_k \searrow 0$ ,  $t_k r_k \to 0$ ,  $x + t_k h + \frac{1}{2} t_k^2 r_k \in C$  and

$$Q_C^{x,\varphi}(h) = \lim_{\substack{k \to \infty \\ 11}} \langle \varphi, r_k \rangle.$$

Note that the above implies in particular that  $h \in \mathcal{T}_C(x)$ . From the outer second-order regularity of C in x, we now obtain

$$\lim_{k \to \infty} \operatorname{dist} \left( r_k, \mathcal{T}_C^2(x, h) \right) = 0,$$

i.e., there exists a sequence  $\{\tilde{r}_k\} \subset \mathcal{T}^2_C(x,h)$  with

(13) 
$$Q_C^{x,\varphi}(h) = \lim_{k \to \infty} \langle \varphi, r_k \rangle = \lim_{k \to \infty} \langle \varphi, \tilde{r}_k \rangle \ge \inf_{r \in \mathcal{T}^2_{\alpha}(x,h)} \langle \varphi, r \rangle.$$

If, on the other hand,  $r \in \mathcal{T}_C^2(x,h)$  is arbitrary but fixed, then we know that there are sequences  $\{t_k\} \subset \mathbb{R}^+$ ,  $\{s_k\} \subset X$  with

$$t_k \searrow 0, \quad x + t_k h + \frac{1}{2} t_k^2 (r + s_k) \in C, \quad s_k \to 0,$$

and we obtain from the definition of  $Q_C^{x,\varphi}(h)$  that

$$Q_C^{x,\varphi}(h) \le \inf \left\{ \liminf_{k \to \infty} \langle \varphi, r_k \rangle \, \middle| \, \begin{cases} \{r_k\} \subset X, \{t_k\} \subset \mathbb{R}^+ : t_k \searrow 0, t_k \, r_k \to 0, \\ x + t_k \, h + \frac{1}{2} \, t_k^2 \, r_k \in C \end{cases} \right\}$$
$$\le \liminf_{k \to \infty} \langle \varphi, r + s_k \rangle = \langle \varphi, r \rangle.$$

This yields

$$Q_C^{x,\varphi}(h) \le \inf_{r \in \mathcal{T}_C^2(x,h)} \langle \varphi, r \rangle$$

which, together with (13), proves the claim.

We would like to point out that Lemma 5.3 cannot be obtained as a corollary of Lemma 5.4. The reason for this is that polyhedric sets do not necessarily have to be second-order regular. In fact, we have the following result.

LEMMA 5.5 (Necessary Condition for Second-Order Regularity). Assume that C is outer second-order regular at  $x \in C$ . Then, for all  $h \in \mathcal{T}_C(x)$ , it holds  $\mathcal{T}_C^2(x,h) \neq \emptyset$  and there exists a positive sequence  $t_k \searrow 0$  such that  $\operatorname{dist}(x + t_k h, C) = \mathcal{O}(t_k^2)$  as  $k \to \infty$ .

*Proof.* Let  $h \in \mathcal{T}_C(x)$  be given. As explained after Definition 5.2, the outer second-order regularity of C at  $x \in C$  implies  $\mathcal{T}_C^2(x,h) \neq \emptyset$ .

Now, let  $r \in \mathcal{T}_C^2(x,h)$  be given. By definition, there is a positive sequence  $t_k \searrow 0$  such that  $\operatorname{dist}(x + t_k h + \frac{1}{2} t_k^2 r, C) = o(t_k^2)$ . Using the latter and the triangle inequality, we obtain

$$\operatorname{dist}(x + t_k h, C) \leq \operatorname{dist}\left(x + t_k h + \frac{1}{2}t_k^2 r, C\right) + \frac{1}{2}t_k^2 \|r\|_X = o(t_k^2) + \mathcal{O}(t_k^2) = \mathcal{O}(t_k^2).$$

This finishes the proof.

Using Lemma 5.5, we can prove that even the most elementary examples of infinite-dimensional polyhedric sets can lack the property of second-order regularity.

Example 5.6. Let (0,1) be equipped with Lebesgue's measure. Define  $X := L^2(0,1)$  and  $C := \{v \in L^2(0,1) : v \geq 0\}$ , and let x be the unique element of X with  $x(\xi) = 1$  for a.a.  $\xi \in (0,1)$ . Then, C is polyhedric at x and it holds  $\mathcal{T}_C(x) = L^2(0,1)$ .

Consider now an arbitrary but fixed  $\alpha \in (-1/2, -1/4)$  and let  $h \in L^2(0,1)$  be defined via  $h(\xi) = -\xi^{\alpha}$  for a.a.  $\xi \in (0,1)$ . Then, for all sequences  $t_k \searrow 0$ , it holds

$$\operatorname{dist}(x + t_k h, C) = \left( \int_0^{t_k^{-1/\alpha}} (1 - t_k \xi^{\alpha})^2 \, \mathrm{d}\xi \right)^{1/2} = c_{\alpha}^{1/2} t_k^{-1/(2\alpha)} \neq \mathcal{O}(t_k^2),$$

where  $c_{\alpha} = \frac{2\alpha^2}{(\alpha+1)(2\alpha+1)} > 0$ . Hence,  $\mathcal{T}_C^2(x,h) = \emptyset$  and, consequently, C cannot be outer second-order regular at x by Lemma 5.5. Using similar arguments, we can show that C fails to be outer second-order regular at all  $x \in C \setminus \{0\}$ .

Note that, due to the effects appearing in Example 5.6, the concept of secondorder regularity is typically not suited for the analysis of optimal control problems with pointwise control or state constraints. It is, however, quite useful when the optimization problem at hand is finite-dimensional or involves constraints of the form  $G(x) \in K$ , where  $K \subset \mathbb{R}^d$  is a closed, convex, non-empty set, cf. Example 5.13. To simplify the derivation of second-order optimality conditions for problems of the latter type, we provide a calculus rule for the curvature of preimages.

Lemma 5.7. Let Z be a Banach space. Assume that  $C = G^{-1}(K)$  holds for some twice continuously Fréchet differentiable function  $G: X \to Z$  and some closed, convex, non-empty set  $K \subset Z$ . Suppose further that a tuple  $(x, \varphi) \in C \times -\mathcal{N}_C^*(x)$  is given such that the Zowe-Kurcyusz constraint qualification

(ZKCQ) 
$$G'(x) X - \mathcal{R}_K(G(x)) = Z$$

is satisfied in x, such that K is second-order regular in G(x), and such that the maps  $h \mapsto G'(x)h$  and  $h \mapsto G''(x)h^2$  are weak-\*-to-strong continuous. Then, C satisfies (MRC) in  $(x,\varphi)$ , it holds  $\mathcal{K}_C^*(x,\varphi) = \varphi^{\perp} \cap G'(x)^{-1}\mathcal{T}_K(G(x))$ , and for every  $h \in \mathcal{K}_C^*(x,\varphi)$  it is true that

$$Q_C^{x,\varphi}(h) = \inf_{r \in G'(x)^{-1} \left(\mathcal{T}_K^2(G(x), G'(x)h) - G''(x)h^2\right)} \langle \varphi, r \rangle.$$

*Proof.* The proof of Lemma 5.7 follows the lines of that of [4, Proposition 3.88]: Let  $h \in \mathcal{K}_C^{\star}(x,\varphi)$  be fixed, and let  $\{r_k\} \subset X, \{t_k\} \subset \mathbb{R}^+$  be sequences satisfying  $t_k \searrow 0$ ,  $t_k r_k \stackrel{\star}{\rightharpoonup} 0$  and  $x + t_k h + \frac{1}{2} t_k^2 r_k \in C$ , i.e.,  $G(x + t_k h + \frac{1}{2} t_k^2 r_k) \in K$ . Then, by a Taylor expansion of G, cf. [6, Theorem 5.6.3], and using our assumptions on G''(x), we obtain

(15) 
$$G\left(x + t_{k} h + \frac{1}{2} t_{k}^{2} r_{k}\right)$$

$$= G(x) + t_{k} G'(x) h + \frac{1}{2} t_{k}^{2} \left(G'(x) r_{k} + G''(x) \left(h + \frac{1}{2} t_{k} r_{k}\right)^{2}\right) + o(t_{k}^{2})$$

$$= G(x) + t_{k} G'(x) h + \frac{1}{2} t_{k}^{2} \left(G'(x) r_{k} + G''(x) h^{2} + \varphi_{k}\right) \in K$$

with some  $\varphi_k$  satisfying  $\varphi_k \to 0$  in Z. Since G'(x) is weak-\*-to-strong continuous, (15) implies  $G'(x)h \in \mathcal{T}_K(G(x))$  and, by [4, Corollary 2.91],  $h \in \mathcal{T}_C(x) = G'(x)^{-1}\mathcal{T}_K(G(x))$ . This yields  $\mathcal{K}_C^{\star}(x,\varphi) = \varphi^{\perp} \cap G'(x)^{-1}\mathcal{T}_K(G(x))$  and shows that it makes sense to use the second-order tangent sets  $\mathcal{T}_C^2(x,h)$  and  $\mathcal{T}_K^2(G(x),G'(x)h)$  in the following. Next, we will prove that  $\operatorname{dist}(r_k,\mathcal{T}_C^2(x,h)) \to 0$ . We start by observing that

$$0 \le D_k := \operatorname{dist} \left( G'(x) r_k, \mathcal{T}_K^2(G(x), G'(x)h) - G''(x)h^2 \right)$$
  
 
$$\le \operatorname{dist} \left( G'(x) r_k + G''(x)h^2 + \varphi_k, \mathcal{T}_K^2(G(x), G'(x)h) \right) + \|\varphi_k\|_Z \to 0,$$

where we used (15),  $t_kG'(x)r_k \to 0$  and the outer second-order regularity of K at G(x).

Now, let  $\{\eta_k\} \subset Z$  be a sequence with  $G'(x)r_k + G''(x)h^2 + \eta_k \in \mathcal{T}_K^2(G(x), G'(x)h)$  and  $\|\eta_k\|_Z \leq D_k + 1/k$ . From [24, Theorem 2.1] and [4, Proposition 2.95], we obtain that there exists a  $\rho > 0$  with

$$B_{\varrho}^{Z}(0) \subset G'(x)B_{1}^{X}(0) - (K - G(x)) \cap B_{1}^{Z}(0).$$

In particular, we may find sequences  $\mu_k \in X$  and  $\lambda_k \in \mathcal{R}_K(G(x))$  such that

(16) 
$$\eta_k = G'(x)\mu_k - \lambda_k \text{ and } \|\mu_k\|_X \le \rho^{-1} \|\eta_k\|_Z.$$

From (16) and the inclusions

(17a) 
$$\mathcal{R}_K(G(x)) \subset \mathcal{T}_{\mathcal{T}_K(G(x))}(h),$$

(17b) 
$$\mathcal{T}_K^2(G(x), G'(x)h) + \mathcal{T}_{\mathcal{T}_K(G(x))}(G'(x)h) \subset \mathcal{T}_K^2(G(x), G'(x)h),$$

which follow from the fact that  $\mathcal{T}_K(G(x))$  is a closed convex cone and [4, Proposition 3.34], we obtain

$$G'(x)(r_k + \mu_k) \in \mathcal{T}_K^2(G(x), G'(x)h) + \lambda_k - G''(x)h^2 \subset \mathcal{T}_K^2(G(x), G'(x)h) - G''(x)h^2.$$

Using the identity

(18) 
$$\mathcal{T}_C^2(x,h) = G'(x)^{-1} \left( \mathcal{T}_K^2(G(x), G'(x)h) - G''(x)(h,h) \right) \quad \forall h \in \mathcal{T}_C(x)$$

that is found, e.g., in [4, Proposition 3.33], we infer

$$r_k + \mu_k \in G'(x)^{-1} \left( \mathcal{T}_K^2(G(x), G'(x)h) - G''(x)h^2 \right) = \mathcal{T}_C^2(x, h).$$

The above implies that we indeed have

$$\operatorname{dist}(r_k, \mathcal{T}_C^2(x, h)) \le \|\mu_k\|_X \le \rho^{-1} \|\eta_k\|_Z \le \rho^{-1} (D_k + 1/k) \to 0.$$

Arguing as in the first part of the proof of Lemma 5.4, we now obtain

$$\liminf_{k \to \infty} \langle \varphi, r_k \rangle \ge \inf_{r \in \mathcal{T}_{\alpha}^2(x,h)} \langle \varphi, r \rangle,$$

and, as a consequence,

$$Q_C^{x,\varphi}(h) \ge \inf_{r \in \mathcal{T}_C^2(x,h)} \langle \varphi, r \rangle.$$

On the other hand, an argumentation analogous to that employed in the second part of the proof of Lemma 5.4 yields

$$Q_C^{x,\varphi}(h) \le \inf \left\{ \liminf_{k \to \infty} \langle \varphi, r_k \rangle \, \middle| \, \begin{cases} \{r_k\} \subset X, \{t_k\} \subset \mathbb{R}^+ : t_k \searrow 0, t_k \, r_k \to 0, \\ x + t_k \, h + \frac{1}{2} \, t_k^2 \, r_k \in C \end{cases} \right\}$$

$$\le \inf_{r \in \mathcal{T}_c^2(x,h)} \langle \varphi, r \rangle \le Q_C^{x,\varphi}(h).$$

Hence, equality holds everywhere, (14) is valid (cf. (18)) and (MRC) is satisfied in  $(x, \varphi)$  by the observation in Remark 3.5. This completes the proof.

Under stronger assumptions on x and G, the right-hand side of (14) is directly related to the directional curvature functional of K.

LEMMA 5.8. In the situation of Lemma 5.7, assume that Z is the dual of a separable Banach space and that there exists a  $\lambda \in \mathcal{N}_K^{\star}(G(x))$  with  $\varphi + G'(x)^{\star}\lambda = 0$  such that (MRC) of K holds in  $(G(x), -\lambda)$  and

(19) 
$$Z = G'(x)X - \mathcal{R}_K(G(x)) \cap \lambda^{\perp}.$$

Then.

(20) 
$$Q_C^{x,\varphi}(h) = Q_K^{G(x),-\lambda}(G'(x)h) + \langle \lambda, G''(x)h^2 \rangle \quad \forall h \in \mathcal{K}_C^{\star}(x,\varphi).$$

*Proof.* Let  $h \in \mathcal{K}_C^{\star}(x,\varphi)$  be arbitrary but fixed. From (17), (19) and [24, Theorem 2.1], it follows (analogously to the proof of Lemma 5.7) that for every  $w \in \mathcal{T}_K^2(G(x), G'(x) h) - G''(x) h^2$  there exist an  $r \in X$  and an  $\eta \in \mathcal{R}_K(G(x)) \cap \lambda^{\perp}$  with

$$w = G'(x) r - \eta$$
 and  $G'(x) r = w + \eta \in \mathcal{T}_K^2(G(x), G'(x) h) - G''(x) h^2$ .

Note that an r with the latter properties necessarily satisfies  $\langle -\lambda, w \rangle = \langle -\lambda, G'(x)r \rangle$  and  $G'(x)r \in G'(x)X \cap (\mathcal{T}_K^2(G(x), G'(x)h) - G''(x)h^2)$ . Consequently, we may deduce

(21)

$$\inf_{w \in \mathcal{T}_K^2(G(x), G'(x) | h) - G''(x)h^2} \langle -\lambda, w \rangle \ge \inf_{w \in G'(x)X \cap \left(\mathcal{T}_K^2(G(x), G'(x) | h) - G''(x)h^2\right)} \langle -\lambda, w \rangle.$$

On the other hand, we trivially have

$$\inf_{w \in \mathcal{T}_K^2(G(x), G'(x) | h) - G''(x)h^2} \langle -\lambda, w \rangle \leq \inf_{w \in G'(x) X \cap \left(\mathcal{T}_K^2(G(x), G'(x) | h) - G''(x)h^2\right)} \langle -\lambda, w \rangle,$$

so equality has to hold in (21). Using this equality, (18), Lemmas 5.4 and 5.7, the identity  $\varphi + G'(x)^*\lambda = 0$  and a straightforward calculation, we obtain

$$\begin{split} & \mathcal{Q}_{C}^{x,\varphi}(h) \\ & = \inf_{r \in \mathcal{T}_{C}^{2}(x,h)} \left\langle \varphi, r \right\rangle = \inf_{r \in G'(x)^{-1} \left( \mathcal{T}_{K}^{2}(G(x), G'(x)h) - G''(x)h^{2} \right)} \left\langle -\lambda, G'(x)r \right\rangle \\ & = \inf_{w \in G'(x)X \cap \left( \mathcal{T}_{K}^{2}(G(x), G'(x)h) - G''(x)h^{2} \right)} \left\langle -\lambda, w \right\rangle = \inf_{w \in \mathcal{T}_{K}^{2}(G(x), G'(x)h) - G''(x)h^{2}} \left\langle -\lambda, w \right\rangle \\ & = \inf_{w \in \mathcal{T}_{K}^{2}(G(x), G'(x)h)} \left\langle -\lambda, w \right\rangle + \left\langle \lambda, G''(x)h^{2} \right\rangle = Q_{K}^{G(x), -\lambda} (G'(x)h) + \left\langle \lambda, G''(x)h^{2} \right\rangle. \end{split}$$

This proves the claim.

Several things are noteworthy regarding Lemma 5.8 and its assumptions. REMARK 5.9.

(i) The pull-back formula (20) is, in fact, valid in a setting that is far more general than the one considered in Lemmas 5.7 and 5.8. It holds, e.g., also for polyhedric sets K provided the strengthened Zowe-Kurcyusz condition (19) is satisfied, cf. [23, Theorem 5.7]. It is further remarkable that the estimate

$$Q_C^{x,\varphi}(h) \ge Q_K^{G(x),-\lambda}(G'(x)h) + \langle \lambda, G''(x)h^2 \rangle \quad \forall h \in \mathcal{K}_C^{\star}(x,\varphi),$$

which yields an SSC for the problem (P), can often be proved without any constraint qualifications at all. To avoid overloading this paper, we leave a detailed discussion of the latter topics for future research.

- (ii) A possible interpretation of the formula (20) is that the (directional) curvature of the set C has its origin in the nonlinearity of G or in the curvedness of K.
- (iii) The condition (19) is well-known and appears, e.g., also in the study of the uniqueness of Lagrange multipliers. It is precisely the ordinary Zowe-Kurcyusz constraint qualification for the set  $\tilde{K} := \{u \in K : (u G(x)) \in \lambda^{\perp}\}$ . We refer to [22, Theorem 2.2] for details on this topic.
- **5.3.** Two Corollaries of Theorem 4.5 and Some Tangible Examples. If we combine the findings of Subsections 5.1 and 5.2 with the analysis of Section 4, then we arrive, e.g., at the following two results.

THEOREM 5.10 (No-Gap Second-Order Condition for Polyhedric Sets). Suppose that X is reflexive and that C is polyhedric at  $\bar{x}$ . Assume that  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$  holds and that  $J''(\bar{x})$  is a Legendre form in the sense of Lemma 5.1 (ii). Then, the condition

$$J''(\bar{x}) h^2 > 0 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{x}, J'(\bar{x})) \setminus \{0\}$$

П

is equivalent to the quadratic growth condition (6) with constants c > 0 and  $\varepsilon > 0$ .

Theorem 5.11 (No-Gap Second-Order Condition under Second-Order Regularity). Let Z be a Banach space. Assume that  $C = G^{-1}(K)$  holds for some twice continuously Fréchet differentiable function  $G: X \to Z$  and some closed, convex, nonempty set  $K \subset Z$ . Assume further that  $J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$  holds, that  $J''(\bar{x})$  is weak- $\star$ lower semicontinuous, that (NDC) holds, that the maps  $h \mapsto G'(\bar{x})h$  and  $h \mapsto G''(\bar{x})h^2$  are weak- $\star$ -to-strong continuous, that K is second-order regular in  $G(\bar{x})$ , and that the constraint qualification

$$G'(\bar{x}) X - \mathcal{R}_K(G(\bar{x})) = Z$$

is satisfied. Then, the condition

(22) 
$$J''(\bar{x}) h^{2} + \inf_{r \in G'(\bar{x})^{-1} \left(\mathcal{T}_{K}^{2}(G(\bar{x}), G'(\bar{x})h) - G''(\bar{x})h^{2}\right)} \langle J'(\bar{x}), r \rangle > 0 \\ \forall h \in J'(\bar{x})^{\perp} \cap G'(\bar{x})^{-1} \mathcal{T}_{K}(G(\bar{x})) \setminus \{0\}$$

is equivalent to the quadratic growth condition (6) with constants c>0 and  $\varepsilon>0$ . If, moreover, we know that Z is the dual of a separable Banach space and that there exists a Lagrange multiplier  $\lambda\in\mathcal{N}_K^\star(G(\bar{x}))$  satisfying  $J'(\bar{x})+G'(\bar{x})^\star\lambda=0$  such that (MRC) holds in  $(G(\bar{x}),-\lambda)$  and such that

$$Z = G'(\bar{x})X - \mathcal{R}_K(G(\bar{x})) \cap \lambda^{\perp},$$

then (22) is equivalent to

$$\partial_{xx}L(x,\lambda)h^2 + Q_K^{G(\bar{x}),-\lambda}(G'(\bar{x})h) > 0 \qquad \forall h \in J'(\bar{x})^{\perp} \cap G'(\bar{x})^{-1}\mathcal{T}_K(G(\bar{x})) \setminus \{0\},$$

where  $L(x,\lambda) := J(x) + \langle \lambda, G(x) \rangle$  is the Lagrangian associated with (P).

We remark that no-gap second-order conditions for specific classes of optimization problems with polyhedric admissible sets (in particular, optimal control problems with box constraints) can be found frequently in the literature. We only mention [2, Theorem 2.7], [8, Theorems 2.2, 2.3], and [9, Theorem 4.13] as examples here. Theorem 5.10 reproduces these results on an abstract level.

Second-order conditions similar to those in Theorem 5.11, on the other hand, have been studied extensively in [4] in various formats and settings, see ibidem Theorems 3.45, 3.83, 3.86, 3.109, 3.137, 3.145, 3.148, 3.155 and Proposition 3.46. It should be noted that the no-gap conditions derived in [4, Section 3.3.3] all require X to be finite-dimensional and, in addition, all need further assumptions on, e.g., the second-order tangent set  $\mathcal{T}_K^2(G(\bar{x}), G'(\bar{x})h)$ . Such assumptions are not needed for the derivation of our second-order condition (22), but may be required for reformulations of (22) as we have seen in the second part of Theorem 5.11.

We conclude this section with two simple examples that demonstrate the usefulness of Theorems 5.10 and 5.11.

Example 5.12 (A Simple Optimal Control Problem with Control Constraints). Consider a minimization problem of the form

(23) 
$$\begin{aligned} & \textit{Minimize} \quad j(y) + \frac{\gamma}{2} \int_{\Omega} u^2 \, \mathrm{d}\mathcal{L}^d \\ & \textit{such that} \quad u \in L^2(\Omega), \quad -1 \leq u \leq 1 \ \textit{a.e. in } \Omega, \quad S(u) = y, \end{aligned}$$

where  $\Omega \subset \mathbb{R}^d$  is a bounded domain,  $j:L^\infty(\Omega) \to \mathbb{R}$  is twice continuously differentiable,  $\gamma > 0$  is a Tikhonov parameter,  $\mathcal{L}^d$  is the Lebesgue measure, and  $S:L^2(\Omega) \to L^\infty(\Omega)$  is (for simplicity) linear and compact. In this situation, the space  $X:=L^2(\Omega)$  is obviously reflexive, the set  $C:=L^2(\Omega,[-1,1])$  is closed, non-empty and polyhedric at every point, and the reduced objective  $J(u):=j(Su)+\frac{\gamma}{2}\int_{\Omega}u^2\,\mathrm{d}\mathcal{L}^d$  is a  $C^2$ -function with

$$J'(u)h = j'(Su)(Sh) + \gamma(u,h)_{L^2(\Omega)} \qquad \forall u,h \in L^2(\Omega),$$
  
$$J''(u)(h_1,h_2) = j''(Su)(Sh_1,Sh_2) + \gamma(h_1,h_2)_{L^2(\Omega)} \quad \forall u,h_1,h_2 \in L^2(\Omega).$$

Note that the map  $h \mapsto J''(u) h^2$  is weakly lower semicontinuous for all  $u \in L^2(\Omega)$ , and that J''(u) is a Legendre form for all  $u \in L^2(\Omega)$  since

$$(24) \quad h_k \rightharpoonup h \text{ and } J''(u) h_k^2 \to J''(u) h^2 \quad \Rightarrow \quad h_k \rightharpoonup h \text{ and } \|h_k\|_{L^2(\Omega)}^2 \to \|h\|_{L^2(\Omega)}^2$$

$$\Rightarrow \quad h_k \to h.$$

Consequently, Theorem 5.10 is applicable in case of problem (23), and we may deduce that, given some  $\bar{u} \in C$  with  $\bar{p} + \gamma \bar{u} \in -\mathcal{N}_C^{\star}(\bar{u})$ , where  $\bar{p} = S^{\star}(j'(S\bar{u}))$  is the adjoint state, the condition

$$j''(S\bar{u})(Sh,Sh) + \gamma ||h||_{L^2(\Omega)}^2 > 0 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{u},\bar{p} + \gamma \bar{u}) \setminus \{0\}$$

is equivalent to the quadratic growth condition

$$J(u) \geq J(\bar{u}) + \frac{c}{2} \|u - \bar{u}\|_{L^2(\Omega)}^2 \qquad \forall u \in C \cap B_{\varepsilon}^{L^2(\Omega)}(\bar{u})$$

with c > 0 and some  $\varepsilon > 0$ . Note that the Tikhonov regularization is of particular importance in the above setting: The condition  $\gamma > 0$  ensures (in combination with the compactness of S) that the derivative J''(u) is a Legendre form for all  $u \in L^2(\Omega)$  and thus guarantees (NDC). We will see in Subsection 6.2 that the situation changes drastically when the regularization parameter  $\gamma$  equals zero.

Example 5.13 (A Simple Optimal Control Problem with a Scalar Constraint). Consider a minimization problem of the form

Minimize 
$$j(y) + \frac{\gamma}{2} \int_{\Omega} u^2 d\mathcal{L}^d$$
  
such that  $u \in L^2(\Omega)$ ,  $Su = y$ ,  $Tu \in B_1^H(0)$ ,

where  $\Omega \subset \mathbb{R}^d$  is a bounded domain,  $j: L^\infty(\Omega) \to \mathbb{R}$  is twice continuously differentiable,  $\gamma > 0$  is a Tikhonov parameter,  $\mathcal{L}^d$  is the Lebesgue measure, H is some Hilbert space, and  $S: L^2(\Omega) \to L^\infty(\Omega)$ ,  $T: L^2(\Omega) \to H$  are (for simplicity) linear and compact. Define  $X:=L^2(\Omega)$ ,  $Z:=\mathbb{R}$ ,  $G(u):=\|Tu\|_H^2-1$ ,  $K:=(-\infty,0]$ ,  $C:=G^{-1}(K)$ , and  $J(u):=j(Su)+\frac{\gamma}{2}\int_{\Omega}u^2\,\mathrm{d}\mathcal{L}^d$ . Then X and Z are Hilbert spaces, C is non-empty, convex and closed, K is non-empty, convex and closed, G is twice continuously differentiable with  $G'(u)h=2(Tu,Th)_H$  and  $G''(u)(h_1,h_2)=2(Th_1,Th_2)_H$  for all  $u\in L^2(\Omega)$ , J is twice continuously differentiable with the same derivatives as in Example 5.12, the map  $h\mapsto J''(u)h^2$  is weakly lower semicontinuous for all  $u\in L^2(\Omega)$ , J''(u) is a Legendre form for all  $u\in L^2(\Omega)$  (cf. Example 5.12), (ZKCQ) is satisfied in every  $u\in C$  (just use a distinction of cases), and K is second-order regular at every  $z\in K$  with

$$\mathcal{T}_{K}(z) = \begin{cases} \mathbb{R} & \text{if } z \in (-\infty, 0), \\ (-\infty, 0] & \text{if } z = 0, \end{cases} \quad \mathcal{T}_{K}^{2}(z, h) = \begin{cases} \mathbb{R} & \text{if } z \in (-\infty, 0), h \in \mathbb{R}, \\ \mathbb{R} & \text{if } z = 0, h \in (-\infty, 0), \\ (-\infty, 0] & \text{if } z = 0, h = 0. \end{cases}$$

Now, let  $\bar{u} \in C$  be given, such that  $\bar{p} + \gamma \bar{u} \in -\mathcal{N}_C^{\star}(\bar{u})$ , where  $\bar{p} = S^{\star}(j'(S\bar{u}))$  is the adjoint state. Using the above observations and (22), we obtain that the condition (25)

$$j''(S\bar{u})(Sh, Sh) + \gamma \|h\|_{L^{2}(\Omega)}^{2} + \inf_{r \in G'(\bar{u})^{-1}\left(\mathcal{T}_{K}^{2}(G(\bar{u}), G'(\bar{u})h) - 2\|Th\|_{H}^{2}\right)} \langle \bar{p} + \gamma \bar{u}, r \rangle > 0$$

$$\forall h \in \left(\bar{p} + \gamma \bar{u}\right)^{\perp} \cap G'(\bar{u})^{-1} \mathcal{T}_{K}(G(\bar{u})) \setminus \{0\}$$

is equivalent to the quadratic growth condition (6) with constants c > 0 and  $\varepsilon > 0$ . Note that  $G'(\bar{u}): L^2(\Omega) \to \mathbb{R}$  is surjective if  $||T\bar{u}||_H > 0$ . Consequently, if  $0 < ||T\bar{u}|| \le 1$ , we may use the second part of Theorem 5.11 to simplify (25). This yields

$$j''(S\bar{u})(Sh, Sh) + \gamma ||h||_{L^{2}(\Omega)}^{2} > 0 \quad \forall h \in (\bar{p} + \gamma \bar{u})^{\perp} \setminus \{0\}$$

for the case  $0 < ||T\bar{u}|| < 1$  and the condition

$$j''(S\bar{u})(Sh, Sh) + \gamma ||h||_{L^{2}(\Omega)}^{2} + 2\lambda ||Th||_{H}^{2} > 0$$

$$\forall h \in (\bar{p} + \gamma \bar{u})^{\perp} \cap \{h \in L^{2}(\Omega) \setminus \{0\} : (Tu, Th)_{H} \leq 0\}$$

for the case  $||T\bar{u}||_H = 1$ . Here,  $\lambda \geq 0$  is the (in this case necessarily unique) Lagrange multiplier associated with  $\bar{u}$ .

We remark that Example 5.13 can also be studied with different means. We chose the approach with the second-order regularity here to illustrate Theorem 5.11.

**6.** Advantages of our Approach. Having demonstrated that the framework of Section 4 indeed allows to reproduce classical results for minimization problems with polyhedric and second-order regular sets, we now turn our attention to the benefits

offered by our approach in comparison with the classical theory. The main advantages of our method are the following.

- (i) Our approach splits the task of proving no-gap second-order optimality conditions for problems of the type (P) into subproblems that can be tackled independently from each other (namely, verifying (NDC), checking the differentiability of J, and computing the directional curvature functional  $Q_C^{x,\varphi}(\cdot)$ ). We can further state our second-order conditions without imposing any preliminary assumptions (as, e.g., polyhedricity or second-order regularity) on the admissible set C, cf. Theorem 4.5. All of this makes our method more flexible than the classical "all-at-once" approach.
- (ii) Our results can also be employed in situations where the admissible set exhibits a singular or degenerate curvature behavior, cf. the examples in Subsections 6.1 and 6.2.
- (iii) Our approach does not require a detailed analysis of the curvature of the set C. To obtain the second-order condition in Theorem 4.5, we only have to study the behavior of the quantity  $\langle J'(\bar{x}), r_k \rangle$  that appears in the definition of the functional  $Q_C^{\bar{x},J'(\bar{x})}(\cdot)$ , i.e., we only have to analyze how the derivative  $J'(\bar{x})$  acts on the second-order corrections  $r_k$  and not how the  $r_k$  behave in detail. This is a major difference to the concept of second-order regularity, cf. Definition 5.2 (iii), and often very advantageous since it allows to exploit additional information about the gradient of the objective. We will see this effect in Subsection 6.2 below.

In the following, we demonstrate by means of two tangible examples that the above points are not only of academic interest but also of relevance in practice. We begin with a simple finite-dimensional optimization problem whose admissible set exhibits a singular curvature behavior.

**6.1. Singular Curvature in Finite Dimensions.** Consider a two-dimensional optimization problem of the form

(26) Minimize 
$$J(x)$$
, such that  $x \in C = \{(x_1, x_2) \in \mathbb{R}^2 \mid x_2 \ge |x_1|^{\alpha}\}$ 

with a twice continuously differentiable objective  $J: \mathbb{R}^2 \to \mathbb{R}$  and some  $\alpha \in (1,2)$ . Set  $\bar{x}:=(0,0)$  and suppose that  $\bar{x}$  is a critical point of (26) with a non-vanishing gradient, i.e.,  $J'(\bar{x})=(0,\beta)\in -\mathcal{N}_C^\star(\bar{x})=\{0\}\times[0,\infty)$  for some  $\beta>0$ . Then, for every critical direction  $h\in \mathcal{K}_C^\star(\bar{x},J'(\bar{x}))=\{(h_1,h_2)\in \mathbb{R}^2\mid h_2=0\}$  and all sequences  $\{t_k\}\subset \mathbb{R}^+$  and  $\{r_k\}\subset \mathbb{R}^2$  satisfying  $t_k\searrow 0$ ,  $t_k\,r_k\to 0$  and  $\bar{x}+t_k\,h+\frac{1}{2}\,t_k^2\,r_k\in C$ , it holds

$$\lim_{k \to \infty} \inf \langle J'(\bar{x}), r_k \rangle 
= 2 \lim_{k \to \infty} \inf J'(\bar{x})^{\top} \frac{\bar{x} + t_k h + \frac{1}{2} t_k^2 r_k}{t_k^2} = 2\beta \lim_{k \to \infty} \inf \frac{(\bar{x} + t_k h + \frac{1}{2} t_k^2 r_k)_2}{t_k^2} 
\geq 2\beta \lim_{k \to \infty} \inf \frac{|(\bar{x} + t_k h + \frac{1}{2} t_k^2 r_k)_1|^{\alpha}}{t_k^2} = 2\beta \lim_{k \to \infty} \inf |(h + \frac{1}{2} t_k r_k)_1|^{\alpha} t_k^{\alpha - 2}.$$

The above implies

$$(28) \quad Q_C^{\bar{x}, J'(\bar{x})}(h) = +\infty \qquad \forall h \in \mathcal{K}_C^{\star}(\bar{x}, J'(\bar{x})) \setminus \{0\} = \{h \in \mathbb{R}^2 \mid h_1 \neq 0, h_2 = 0\}.$$

Using (28), Theorem 4.5 and the fact that the conditions (MRC) and (NDC) are trivially satisfied for (26), we obtain (analogous to [4, Example 3.84]) the following result.

THEOREM 6.1. If  $\bar{x} = (0,0)$  is a critical point of (26) with  $J'(\bar{x}) \neq 0$ , then  $\bar{x}$  is a local minimizer of (26) and there exist parameters c > 0 and  $\varepsilon > 0$  such that the quadratic growth condition (6) is satisfied.

Note that the second derivative  $J''(\bar{x})$  is not important for local optimality of  $\bar{x}$ . The reason for this is that the curvature of the boundary  $\partial C$  is singular at the origin and can thus compensate for any negative curvature that the objective J might have at  $\bar{x}$ . It should be noted further that the set C in (26) is neither polyhedric (trivially) nor second-order regular at  $\bar{x}$  (since  $\mathcal{T}_C^2(\bar{x},h)=\emptyset$  for all  $h\in\mathcal{K}_C^{\star}(\bar{x},J'(\bar{x}))\setminus\{0\}$ , cf. Lemma 5.5, and the estimate (27)). This demonstrates that (26) does not fall under the setting of Theorems 5.10 and 5.11 and is indeed not covered by what is typically seen as the classical second-order theory.

We remark that, given an optimization problem of the form

(29) Minimize 
$$J(x)$$
, such that  $x \in C = \{(x_1, x_2) \in \mathbb{R}^2 \mid x_2 \le |x_1|^{\alpha} \}$ 

with  $J \in C^2(\mathbb{R}^2)$ ,  $J'(\bar{x}) = (0, -\beta) \in -\mathcal{N}_C^{\star}(\bar{x}) = \{0\} \times (-\infty, 0]$ ,  $\beta > 0$ ,  $\bar{x} := (0, 0)$ , we can use exactly the same arguments as for (26) to prove

$$Q_C^{\bar{x},J'(\bar{x})}(h) = -\infty \qquad \forall h \in \mathcal{K}_C^{\star}(\bar{x},J'(\bar{x})) = \{h \in \mathbb{R}^2 \mid h_2 = 0\}.$$

The above yields in combination with Theorems 4.2 and 4.3 that  $\bar{x} = (0,0)$  can never be a local minimizer of (29) unless the derivative  $J'(\bar{x})$  is identical zero.

**6.2.** No-Gap Second-Order Conditions for Bang-Bang Problems. In this section, we demonstrate that the analysis of Section 4 is not only relevant for finite-dimensional toy problems à la (26) and (29), but also applicable in more complicated situations. In what follows, we will use it to derive no-gap second-order conditions for bang-bang optimal control problems. As a motivation, let us consider the optimization problem (23) in Example 5.12 with  $\gamma = 0$ , i.e., the problem

(30) Minimize 
$$j(y)$$
  
such that  $u \in L^2(\Omega)$ ,  $-1 \le u \le 1$  a.e. in  $\Omega$ ,  $S(u) = y$ .

From Theorem 4.2, we obtain that every minimizer  $\bar{u}$  of (30) satisfies  $\bar{p} \in -\mathcal{N}_C^{\star}(\bar{u})$ , where  $\bar{p} = S^{\star}(j'(S\bar{u}))$  is the adjoint state. In particular, this implies that a minimizer  $\bar{u}$  with  $\mathcal{L}^d(\{\bar{p}=0\}) = 0$  can only take the values  $\pm 1$  a.e. in  $\Omega$ . Such a solution  $\bar{u}$  is called bang-bang.

The major problem that arises when SSC for a bang-bang control  $\bar{u}$  are considered is the verification of the non-degeneracy condition (NDC). Recall that in Example 5.12 the latter is satisfied since the Tikhonov regularization causes the second derivative of the reduced objective to be a Legendre form in  $L^2(\Omega)$ , cf. (24). For (30) such an argumentation is obviously not possible and this is not an artificial problem: It can be shown that quadratic growth in  $L^2(\Omega)$  is in general not possible for a bangbang solution  $\bar{u}$  of (30), i.e., the growth condition (6) typically does not hold with  $X = L^2(\Omega)$  and  $c, \varepsilon > 0$ , cf. [7, end of Section 2].

Hence, Theorem 4.5 cannot be applicable when we work with the space  $X = L^2(\Omega)$ . Note that, if we calculate the critical cone  $\mathcal{K}_C^{\star}(\bar{u},\bar{p})$  for a bang-bang control  $\bar{u}$  in the  $L^2$ -setting, then we end up with  $\mathcal{K}_C^{\star}(\bar{u},\bar{p}) = \mathcal{T}_C(\bar{u}) \cap \bar{p}^{\perp} = \{0\}$  so that the conditions  $j''(S\bar{u})(Sh,Sh) > 0 \ \forall h \in \mathcal{K}_C^{\star}(\bar{u},\bar{p}) \setminus \{0\}$  and  $j''(S\bar{u})(Sh,Sh) \geq 0 \ \forall h \in \mathcal{K}_C^{\star}(\bar{u},\bar{p})$ , which are the natural candidates for the SSC and SNC, respectively,

are both void. This also indicates that it is not useful to discuss (30) as a problem in  $X = L^2(\Omega)$ .

The above discussion shows that, if we want to derive no-gap second-order conditions for a bang-bang optimal control problem of the type (30), then we have to work with a space X that is different from  $L^2(\Omega)$ . As it turns out, the right choice is the measure space  $X = \mathcal{M}(\Omega)$ . Therefore, we consider the following setting.

ASSUMPTION 6.2 (Standing Assumptions and Notation for the Bang-Bang Setting). We suppose that  $\Omega \subset \mathbb{R}^d$ ,  $d \geq 1$ , is a bounded domain with a Lipschitz boundary, cf. [16, Definition 1.2.1.1]. We define the space  $Y := C_0(\Omega) = \operatorname{cl}_{\|.\|_{\infty}}(C_c(\Omega))$  endowed with the usual supremum norm. Its dual space can be identified with  $X := \mathcal{M}(\Omega)$  which is the space of signed finite Radon measures on  $\Omega$  endowed with the norm  $\|\mu\|_{\mathcal{M}(\Omega)} := |\mu|(\Omega)$ , cf. [1, Theorem 1.54]. The space  $L^1(\Omega)$  is identified with a closed subspace of  $\mathcal{M}(\Omega)$  via the isometric embedding  $x \mapsto x\mathcal{L}^d$ , where  $\mathcal{L}^d$  is Lebesgue's measure. Note that this implies  $\|x\|_{L^1(\Omega)} = \|x\|_{\mathcal{M}(\Omega)}$  for all  $x \in L^1(\Omega)$ . Finally,  $C := L^{\infty}(\Omega, [-1, 1]) = \{x \in L^{\infty}(\Omega) : -1 \leq x \leq 1 \text{ a.e. in } \Omega\} \subset \mathcal{M}(\Omega)$ .

The reason for using the space  $X = \mathcal{M}(\Omega)$  is the following observation, cf. [11, Proposition 2.7].

LEMMA 6.3. Let  $\bar{x} \in C$  and  $\bar{\varphi} \in -\mathcal{N}_C^{\star}(\bar{x})$  be given. Define

(31) 
$$K(\bar{\varphi}) := \frac{1}{4} \liminf_{s \searrow 0} \left( \frac{s}{\mathcal{L}^d(\{|\bar{\varphi}| \le s\})} \right) \in [0, +\infty].$$

Then, there exists a family of constants  $\{c_{\varepsilon}\}$  satisfying  $c_{\varepsilon} \searrow 0$  as  $\varepsilon \searrow 0$  such that

$$(32) \ \langle \bar{\varphi}, x - \bar{x} \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} \ge \left( \frac{1}{2} K(\bar{\varphi}) - c_{\varepsilon} \right) \|x - \bar{x}\|_{L^1(\Omega)}^2 \quad \forall x \in C \cap B_{\varepsilon}^X(\bar{x}) \quad \forall \varepsilon > 0.$$

Proof. We adapt the proof of [11, Proposition 2.7]. Define  $K := K(\bar{\varphi})$ . If K = 0, then the claim is trivially true. If K > 0, then it necessarily holds that  $\mathcal{L}^d(\{\bar{\varphi} = 0\}) = 0$  and we obtain from  $\bar{\varphi} \in -\mathcal{N}_C^{\star}(\bar{x})$  that  $\bar{x}$  is bang-bang with  $\bar{x} = -\operatorname{sgn}\bar{\varphi}$  a.e. in  $\Omega$ . Using the latter and  $\mathcal{N}_C^{\star}(\bar{x}) \subset C_0(\Omega)$ , we may calculate that for all  $h \in X$  and all t > 0 with  $\bar{x} + th \in C$  and  $||h||_{X} = ||h||_{L^1(\Omega)} = 1$ , it holds

$$\int_{\Omega} \bar{\varphi} \frac{h}{t} d\mathcal{L}^{d} = \int_{\Omega} |\bar{\varphi}| \frac{|h|}{t} d\mathcal{L}^{d} \ge \int_{\{|\bar{\varphi}| > Kt\}} |\bar{\varphi}| \frac{|h|}{t} d\mathcal{L}^{d} 
\ge \int_{\Omega} K|h| d\mathcal{L}^{d} - \int_{\{|\bar{\varphi}| < Kt\}} K|h| d\mathcal{L}^{d} \ge K - K||h||_{L^{\infty}(\Omega)} \mathcal{L}^{d}(\{|\bar{\varphi}| \le Kt\}).$$

By using  $t := \|x - \bar{x}\|_{L^1(\Omega)}$ ,  $h := (x - \bar{x})/\|x - \bar{x}\|_{L^1(\Omega)}$  and  $\|h\|_{L^{\infty}(\Omega)} \le 2/t$ , this implies

$$\langle \bar{\varphi}, x - \bar{x} \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} \ge \left( K - 2K^2 \sup_{0 < t < \varepsilon} \frac{\mathcal{L}^d(\{|\bar{\varphi}| \le Kt\})}{Kt} \right) \|x - \bar{x}\|_{L^1(\Omega)}^2$$

for all  $x \in C \cap B_{\varepsilon}^X(\bar{x}) \setminus \{\bar{x}\}$  and all  $\varepsilon > 0$ . Note that the coefficient on the right-hand side of the last estimate satisfies

$$\lim_{\varepsilon \searrow 0} \left( K - 2K^2 \sup_{0 < t < \varepsilon} \frac{\mathcal{L}^d(\{|\bar{\varphi}| \le Kt\})}{Kt} \right) = K - 2K^2 \left( \limsup_{t \searrow 0} \frac{\mathcal{L}^d(\{|\bar{\varphi}| \le t\})}{t} \right) = \frac{1}{2}K.$$

This proves the claim.

Lemma 6.3 shows that Lemma 5.1 (iii) is applicable when we consider a bangbang solution  $\bar{x}$  whose gradient  $\bar{\varphi} := J'(\bar{x})$  satisfies  $K(\bar{\varphi}) > 0$ . This allows us to verify (NDC) and to obtain the following result from Theorem 4.5.

THEOREM 6.4 (No-Gap Second-Order Condition for Bang-Bang Problems). We consider an optimization problem of the form (P) with C, X etc. as in Assumption 6.2. Assume that  $\bar{x} \in C$  is fixed, that J satisfies the conditions in Assumption 4.1, that the map  $X \ni h \mapsto J''(\bar{x}) h^2 \in \mathbb{R}$  is weak-\* continuous, that  $\bar{\varphi} := J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$ , and that the constant  $K(\bar{\varphi})$  in (31) is positive. Then, the condition

$$Q_C^{\bar{x},\bar{\varphi}}(h) + J''(\bar{x})h^2 > 0 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{x},\bar{\varphi}) \setminus \{0\}$$

is equivalent to the quadratic growth condition

(34) 
$$J(x) \ge J(\bar{x}) + \frac{c}{2} \|x - \bar{x}\|_{L^{1}(\Omega)}^{2} \quad \forall x \in C \cap B_{\varepsilon}^{X}(\bar{x})$$

with constants c > 0 and  $\varepsilon > 0$ .

Proof. We only need to check that Theorem 4.5 is applicable. The setting in Assumption 6.2 clearly fits into that of Assumption 2.1, and Assumption 4.1 is trivially satisfied. From  $K(\bar{\varphi}) > 0$  and Lemma 6.3, we further obtain that there exist constants  $c, \varepsilon > 0$  with  $J'(\bar{x})(x - \bar{x}) \geq \frac{c}{2} \|x - \bar{x}\|^2$  for all  $x \in B_{\varepsilon}^X(\bar{x}) \cap C$ . This yields, in combination with the weak-\* continuity of  $h \mapsto J''(\bar{x})h^2$  and Lemma 5.1 (iii), that (NDC) holds. Note that the weak- $\star$  continuity of  $h \mapsto J''(\bar{x})h^2$  also implies (i) in Theorem 4.3. This shows that Theorem 4.5 is applicable and proves the claim.

Some remarks concerning Lemma 6.3 and Theorem 6.4 are in order.

Remark 6.5.

- (i) Theorem 6.4 provides no-gap second-order conditions even in the case that we cannot characterize the directional curvature functional  $Q_C^{\bar{x},\bar{\varphi}}(\cdot)$  precisely.
- Recall that the condition in Lemma 5.1 (iii) not only implies (NDC) but also yields a coercivity estimate for the functional  $Q_C^{\bar{x},\bar{\varphi}}(\cdot)$ , see (12). Using this estimate and (32), we obtain that, in the situation of Theorem 6.4,

(35) 
$$Q_C^{\bar{x},\bar{\varphi}}(h) \ge K(\bar{\varphi}) \|h\|_X^2 \qquad \forall h \in \mathcal{K}_C^{\star}(\bar{x},\bar{\varphi}).$$

Here,  $K(\bar{\varphi}) > 0$  is again defined by (31). We point out that (35) implies that the set  $C = L^{\infty}(\Omega, [-1, 1])$  possesses positive curvature as a subset of the space  $\mathcal{M}(\Omega)$ . This is not true if C is considered as a subset of the space  $L^{2}(\Omega)$  as we have seen in Example 5.12 (in  $L^{2}(\Omega)$ , C is polyhedric and the curvature functional is zero).

(iii) From (35), it follows that  $J''(\bar{x}) h^2 > -K(\bar{\varphi}) \|h\|_X^2$  for all  $h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi}) \setminus \{0\}$ is a sufficient condition for quadratic growth in the situation of Theorem 6.4. cf. (33). We point out that this SSC is sharper than that found in [11, Corollary 2.15]. In this contribution the authors work with the slightly more restrictive "global" level set assumption

$$\tilde{K} \leq \frac{s}{4 \, \mathcal{L}^d(\{|\bar{\varphi}| \leq s\})} \quad \forall s > 0$$

for some  $\tilde{K} > 0$ . Note that such a  $\tilde{K}$  necessarily satisfies  $\tilde{K} \leq K(\bar{\varphi})$ . We further point out that the SSC in [11, Corollary 2.15] can be improved to

$$\exists \varepsilon > 0: \quad J''(\bar{x}) \, h^2 \ge -\big(\tilde{K} - \varepsilon\big) \|h\|_X^2 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi}),$$

cf. [10, Theorem 2.4].

(iv) We expect that the SSC  $J''(\bar{x}) h^2 > -K(\bar{\varphi}) \|h\|_X^2 \ \forall h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi}) \setminus \{0\}$  can also be formulated as an inequality on the so-called extended critical cone introduced in [7], cf. [11, Theorem 2.14] and [10, Theorem 2.4]. We do not pursue this approach here.

The next step is the calculation of the curvature functional  $Q_C^{\bar{x},\bar{\varphi}}(\cdot)$  for a bangbang solution  $\bar{x}$  in order to obtain a no-gap optimality condition that is more explicit than (33). Hence, we have to compute the (directional) curvature of the set  $C := L^{\infty}(\Omega, [-1,1])$  as a subset of the space  $\mathcal{M}(\Omega)$ . In the remainder of this section, we will consider a bang-bang solution  $\bar{x}$  whose gradient  $\bar{\varphi} := J'(\bar{x}) \in -\mathcal{N}_C^{\star}(\bar{x})$  is in  $C^1(\Omega)$ . Let us first fix our assumptions on the  $\bar{x}$  under consideration.

Assumption 6.6 (Assumptions and Notation for the Calculation of  $Q_C^{\bar{x},\bar{\varphi}}(\cdot)$ ).

In addition to Assumption 6.2, we suppose that  $\bar{x} \in C$  and  $\bar{\varphi} \in -\mathcal{N}_C^{\star}(\bar{x})$  are given. We require  $\bar{\varphi} \in \iota(C_0(\Omega) \cap C^1(\Omega))$  and define  $\mathcal{Z} := \{z \in \Omega : \bar{\varphi}(z) = 0\}$ . We assume  $\mathcal{Z} \subset \{z \in \Omega : |\nabla \bar{\varphi}(z)| \neq 0\}$ . Here and in the sequel,  $|\nabla \bar{\varphi}(z)|$  denotes the Euclidean norm of  $\nabla \bar{\varphi}(z) \in \mathbb{R}^d$ .

Finally, we denote by  $\mathcal{H}^{d-1}$  the (d-1)-dimensional Hausdorff measure, which is scaled as in [15, Definition 2.1].

In the above situation, the set  $\mathcal{Z}$  is a (d-1)-dimensional  $C^1$ -submanifold of  $\mathbb{R}^d$  due to the implicit function theorem, cf. [17, Theorem 2.32]. This implies in particular that  $\mathcal{L}^d(\mathcal{Z}) = \mathcal{L}^d(\{\bar{\varphi} = 0\}) = 0$  and that  $\bar{x}$  is indeed bang-bang with  $\bar{x} = -\operatorname{sgn}\bar{\varphi}$  a.e. in  $\Omega$ . To calculate  $Q_C^{\bar{x},\bar{\varphi}}(\cdot)$ , we need the following directional Taylor-like expansion of the  $L^1(\Omega)$ -norm.

LEMMA 6.7 ([12, Corollary 5.10]). Given Assumption 6.6, for all  $v \in C_c(\Omega) \cap H^1(\Omega)$  and all sequences  $t_k \in (0, \infty)$  with  $t_k \searrow 0$ , it is true that

(36) 
$$\int_{\Omega} |-\bar{\varphi} + t_k v| d\mathcal{L}^d = \int_{\Omega} |\bar{\varphi}| d\mathcal{L}^d + t_k \int_{\Omega} \bar{x} v d\mathcal{L}^d + t_k^2 \int_{\mathcal{Z}} \frac{v^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} + o(t_k^2).$$

Proof in the case d=1. To give the reader an idea of how Lemma 6.7 is obtained, we prove (36) in the one-dimensional setting. The proof of the general case is similar but much more technical, see [12, Corollary 5.10]. So let us suppose that d=1 and that  $v \in C_c(\Omega) \cap H^1(\Omega)$  and  $\{t_k\} \subset (0,\infty)$  with  $t_k \searrow 0$  are given. Then,  $\Omega$  is an interval and the compactness of the support  $\sup(v)$ , the regularity of  $\bar{\varphi}$  and our assumption  $\mathcal{Z} \subset \{z \in \Omega : |\bar{\varphi}'(z)| \neq 0\}$  yield that the set  $\mathcal{Z} \cap \sup(v)$  is finite. Denote the elements of  $\mathcal{Z} \cap \sup(v)$  with  $a_i$ , i=1,...,n, assume that  $a_1 < a_2 < ... < a_n$  holds and choose  $b_i$ , i=1,...,n+1, such that  $b_i < a_i < b_{i+1}$  for all i=1,...,n and  $\sup(v) \subset [b_1,b_{n+1}] \subset \Omega$ . Then, we may write

$$(37) \frac{1}{t_k} \int_{\Omega} \frac{|-\bar{\varphi} + t_k v| - |\bar{\varphi}|}{t_k} - \bar{x}v \, d\mathcal{L}^1 = \sum_{i=1}^n \frac{1}{t_k} \int_{b_i}^{b_{i+1}} \frac{|-\bar{\varphi} + t_k v| - |\bar{\varphi}|}{t_k} + \operatorname{sgn}(\bar{\varphi})v \, d\mathcal{L}^1.$$

Consider now an arbitrary but fixed  $i \in \{1, ..., n\}$ , assume w.l.o.g. that  $\bar{\varphi}'(a_i) > 0$  (the case  $\bar{\varphi}'(a_i) < 0$  is analogous) and choose an  $\varepsilon > 0$  such that  $\bar{\varphi}' \geq \delta > 0$  holds in  $(a_i - \varepsilon, a_i + \varepsilon) \subset (b_i, b_{i+1})$ . Then, it follows from our construction, the boundedness of v, the fact that  $\bar{\varphi} \leq -c < 0$  and  $0 < c \leq \bar{\varphi}$  holds in  $[b_i, a_i - \varepsilon]$  and  $[a_i + \varepsilon, b_{i+1}]$  for

some c > 0, respectively, and a simple distinction of cases that

$$\frac{1}{t_k} \int_{b_i}^{b_{i+1}} \frac{|-\bar{\varphi} + t_k v| - |\bar{\varphi}|}{t_k} + \operatorname{sgn}(\bar{\varphi}) v \, d\mathcal{L}^1$$

$$= \int_{b_i}^{a_i} 2 \frac{\max(0, \bar{\varphi} - t_k v)}{t_k^2} \, d\mathcal{L}^1 + \int_{a_i}^{b_{i+1}} 2 \frac{\max(0, -\bar{\varphi} + t_k v)}{t_k^2} \, d\mathcal{L}^1$$

$$= \int_{a_i - \varepsilon}^{a_i} 2 \frac{\max(0, \bar{\varphi} - t_k v)}{t_k^2} \, d\mathcal{L}^1 + \int_{a_i}^{a_i + \varepsilon} 2 \frac{\max(0, -\bar{\varphi} + t_k v)}{t_k^2} \, d\mathcal{L}^1 + o(1),$$

where the Landau symbol refers to the limit  $k \to \infty$ . Note that the integrand of the second integral on the right-hand side of (38) is only non-zero in a  $z \in (a_i, a_i + \varepsilon)$  if

$$\delta(z - a_i) \le \int_{a_i}^z \bar{\varphi}'(s) d\mathcal{L}^1(s) = \bar{\varphi}(z) \le t_k v(z) \le t_k ||v||_{L^{\infty}(\Omega)}.$$

Consequently, for all large enough k, we have

$$\int_{a_{i}}^{a_{i}+\varepsilon} 2 \frac{\max(0, -\bar{\varphi} + t_{k}v)}{t_{k}^{2}} d\mathcal{L}^{1}$$

$$= \int_{a_{i}}^{a_{i}+Ct_{k}} 2 \frac{\max(0, -\bar{\varphi}(z) + t_{k}v(z))}{t_{k}^{2}} d\mathcal{L}^{1}(z)$$

$$= \int_{0}^{C} 2 \frac{\max(0, -\bar{\varphi}(a_{i} + t_{k}z) + t_{k}v(a_{i} + t_{k}z))}{t_{k}^{2}} t_{k} d\mathcal{L}^{1}(z)$$

$$= \int_{0}^{C} 2 \max\left(0, -\int_{0}^{1} \bar{\varphi}'(a_{i} + st_{k}z) dsz + v(a_{i} + t_{k}z)\right) d\mathcal{L}^{1}(z)$$

$$= \int_{0}^{C} 2 \max\left(0, -\bar{\varphi}'(a_{i})z + v(a_{i})\right) d\mathcal{L}^{1}(z) + o(1),$$

where  $C := ||v||_{L^{\infty}(\Omega)}/\delta$  and where the last identity follows from the dominated convergence theorem. From  $\bar{\varphi}'(a_i) > 0$ , we now obtain

$$\int_{0}^{C} 2 \max (0, -\bar{\varphi}'(a_{i})z + v(a_{i})) d\mathcal{L}^{1}(z)$$

$$= 2 \int_{0}^{\max(0, v(a_{i}))/\bar{\varphi}'(a_{i})} -\bar{\varphi}'(a_{i})z + v(a_{i}) d\mathcal{L}^{1}(z) = \frac{\max(0, v(a_{i}))^{2}}{\bar{\varphi}'(a_{i})}.$$

If we use exactly the same argumentation for the first integral on the right-hand side of (38) and combine our results with (37), then we arrive at the identity

$$\frac{1}{t_k} \int_{\Omega} \frac{|-\bar{\varphi} + t_k v| - |\bar{\varphi}|}{t_k} - \bar{x}v \, d\mathcal{L}^1 = \sum_{i=1}^n \frac{v(a_i)^2}{|\bar{\varphi}'(a_i)|} + o(1).$$

Rewriting the above yields (36) in the case d=1. This completes the proof.

The next lemma provides a link between the curvature of C and  $\nabla \bar{\varphi}$ .

LEMMA 6.8. For all  $v \in C_c(\Omega)$ , all  $h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$ , and all  $\alpha > 0$ , it holds

(39) 
$$\frac{\alpha^2}{2} Q_C^{\bar{x},\bar{\varphi}}(h) - \alpha \langle v, h \rangle_{C_0(\Omega),\mathcal{M}(\Omega)} + \int_{\mathcal{Z}} \frac{v^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} \ge 0.$$

*Proof.* Let  $\alpha > 0$  and  $h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$  be given. From the cone property of  $\mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$ , it follows  $\alpha h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$ . This implies that there exist sequences  $\{r_k\} \subset X$ ,  $\{t_k\} \subset \mathbb{R}^+$  with  $t_k \searrow 0$ ,  $t_k r_k \stackrel{\star}{\rightharpoonup} 0$  and  $\bar{x} + t_k \alpha h + \frac{1}{2} t_k^2 r_k \in C$ . Fix such sequences  $\{r_k\}$ ,  $\{t_k\}$  and define  $h_k := \alpha h + \frac{1}{2} t_k r_k$ . Then, it holds  $h_k \in \mathcal{R}_C(\bar{x}) \subset L^{\infty}(\Omega)$  and  $h_k \stackrel{\star}{\rightharpoonup} \alpha h$  in X, and Lemma 6.7 yields

$$t_{k}^{2} \int_{\mathcal{Z}} \frac{v^{2}}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} + o(t_{k}^{2})$$

$$= \left(\sup_{q \in L^{\infty}(\Omega, [-1,1])} \int_{\Omega} (-\bar{\varphi} + t_{k}v) q d\mathcal{L}^{d}\right) - \int_{\Omega} (|\bar{\varphi}| + t_{k}\bar{x}v) d\mathcal{L}^{d}$$

$$= \left(\sup_{p \in L^{\infty}(\Omega): \bar{x} + t_{k}p \in L^{\infty}(\Omega, [-1,1])} \int_{\Omega} (-\bar{\varphi} + t_{k}v) (\bar{x} + t_{k}p) d\mathcal{L}^{d}\right) - \int_{\Omega} (|\bar{\varphi}| + t_{k}\bar{x}v) d\mathcal{L}^{d}$$

$$= \left(\sup_{p \in L^{\infty}(\Omega): \bar{x} + t_{k}p \in L^{\infty}(\Omega, [-1,1])} \int_{\Omega} (-\bar{\varphi} + t_{k}v) (t_{k}p) d\mathcal{L}^{d}\right)$$

$$= \left(\sup_{p \in L^{\infty}(\Omega): \bar{x} + t_{k}p \in L^{\infty}(\Omega, [-1,1])} \int_{\Omega} (-\bar{\varphi} + t_{k}v) (t_{k}p) d\mathcal{L}^{d}\right)$$

$$\geq \int_{\Omega} (-\bar{\varphi} + t_{k}v) (t_{k}h_{k}) d\mathcal{L}^{d} \qquad \forall v \in C_{c}(\Omega) \cap H^{1}(\Omega).$$

If we divide (40) by  $t_k^2$  and let  $k \to \infty$ , then we obtain (with  $\langle \bar{\varphi}, h_k \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} = \langle \bar{\varphi}, \frac{1}{2} t_k r_k \rangle_{C_0(\Omega), \mathcal{M}(\Omega)}$ )

$$\frac{1}{2} \liminf_{k \to \infty} \langle \bar{\varphi}, r_k \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} - \langle v, \alpha h \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} + \int_{\mathcal{Z}} \frac{v^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} \ge 0$$

for all  $v \in C_c(\Omega) \cap H^1(\Omega)$ . Taking the infimum over all sequences  $\{r_k\}$ ,  $\{t_k\}$ , using the positive homogeneity of the functional  $Q_C^{\bar{x},\bar{\varphi}}(\cdot)$ , and employing a density argument, we now arrive at

$$\frac{\alpha^2}{2} Q_C^{\bar{x},\bar{\varphi}}(h) - \langle v, \alpha h \rangle_{C_0(\Omega),\mathcal{M}(\Omega)} + \int_{\mathcal{Z}} \frac{v^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} \ge 0 \qquad \forall v \in C_c(\Omega).$$

П

This is the desired estimate.

We are now in the position to prove a lower bound for the curvature of C.

Proposition 6.9. For all  $h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$  with  $Q_C^{\bar{x}, \bar{\varphi}}(h) < \infty$  there exists a g such that

$$h = g\mathcal{H}^{d-1}|_{\mathcal{Z}}, \qquad g \in L^{1}\left(\mathcal{Z}, \mathcal{H}^{d-1}\right) \cap L^{2}\left(\mathcal{Z}, |\nabla \bar{\varphi}| \mathcal{H}^{d-1}\right),$$
$$\frac{1}{2} \int_{\mathcal{Z}} g^{2} |\nabla \bar{\varphi}| d\mathcal{H}^{d-1} \leq Q_{C}^{\bar{x}, \bar{\varphi}}(h).$$

*Proof.* Let  $h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi}) \subset \mathcal{M}(\Omega)$  with  $Q_C^{\bar{x}, \bar{\varphi}}(h) < \infty$  be given. Then, for all  $v \in C_c(\Omega)$  with v = 0 on  $\mathcal{Z}$ , we obtain from (39) that

$$\frac{\alpha}{2} Q_C^{\bar{x},\bar{\varphi}}(h) \ge \langle v, h \rangle_{C_0(\Omega),\mathcal{M}(\Omega)} \quad \forall \alpha > 0,$$

i.e.,  $\langle v, h \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} \leq 0$ . Using  $\pm v$ , we find  $\langle v, h \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} = 0$ . Hence, the map

$$\tilde{h}: C_c(\mathcal{Z}) \to \mathbb{R}, \qquad \tilde{v} \mapsto \langle v, h \rangle_{C_0(\Omega), \mathcal{M}(\Omega)}, \quad v \in C_c(\Omega), \ v|_{\mathcal{Z}} = \tilde{v}$$

is well-defined as it is independent of the extension v of  $\tilde{v}$  appearing in its definition. Note that, given a  $\tilde{v} \in C_c(\mathcal{Z})$ , we can always find a  $v \in C_c(\Omega)$  with  $v|_{\mathcal{Z}} = \tilde{v}$ , cf. the submanifold property of  $\mathcal{Z}$ . From (39) with

$$\alpha = \left( \int_{\mathcal{Z}} \frac{\tilde{v}^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} \right)^{1/2} \beta, \quad \beta > 0 \text{ arbitrary but fixed,}$$

it now follows

$$\left(\frac{\beta}{2}Q_C^{\bar{x},\bar{\varphi}}(h) + \frac{1}{\beta}\right) \left(\int_{\mathcal{Z}} \frac{\tilde{v}^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1}\right)^{1/2} \ge \tilde{h}(\tilde{v}) \quad \forall \tilde{v} \in C_c(\mathcal{Z}).$$

Now, since  $C_c(\mathcal{Z})$  is dense in the Lebesgue-space  $L^2(\mathcal{Z}, \mathcal{H}^{d-1}/|\nabla \bar{\varphi}|)$ , the functional  $\tilde{h}$  can be uniquely extended. Thus, there exists an  $f \in L^2(\mathcal{Z}, \mathcal{H}^{d-1}/|\nabla \bar{\varphi}|)$  with

$$\left(\frac{\beta}{2}Q_C^{\bar{x},\bar{\varphi}}(h) + \frac{1}{\beta}\right) \left(\int_{\mathcal{Z}} \frac{\tilde{v}^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1}\right)^{1/2} \ge \tilde{h}(\tilde{v}) = \int_{\mathcal{Z}} \frac{\tilde{v}f}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} \quad \forall \tilde{v} \in C_c(\mathcal{Z}).$$

Note that f is independent of  $\beta$  due to the density of  $C_c(\mathcal{Z})$  in  $L^2(\mathcal{Z}, \mathcal{H}^{d-1}/|\nabla \bar{\varphi}|)$ . We thus arrive at

$$\langle v, h \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} = \tilde{h}(v|_{\mathcal{Z}}) = \int_{\mathcal{Z}} \frac{vf}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1} \quad \forall v \in C_c(\Omega)$$

with

$$\left(\int_{\mathcal{Z}} \frac{f^2}{|\nabla \bar{\varphi}|} d\mathcal{H}^{d-1}\right)^{1/2} \le \left(\frac{\beta}{2} Q_C^{\bar{x}, \bar{\varphi}}(h) + \frac{1}{\beta}\right) \qquad \forall \beta > 0.$$

Choosing  $\beta = (2/Q_C^{\bar{x},\bar{\varphi}}(h))^{1/2}$  for  $Q_C^{\bar{x},\bar{\varphi}}(h) > 0$  and  $\beta$  arbitrarily large for  $Q_C^{\bar{x},\bar{\varphi}}(h) = 0$ , defining  $g := f/|\nabla \bar{\varphi}|$  and using the density of  $C_c(\Omega)$  in  $C_0(\Omega)$  now yields the claim. Note that  $g \in L^1(\mathcal{Z},\mathcal{H}^{d-1})$  follows trivially from  $h = g\mathcal{H}^{d-1}|_{\mathcal{Z}} \in \mathcal{M}(\Omega)$ .

Next, we address the reverse estimate to that in Proposition 6.9.

LEMMA 6.10. Let  $g \in C_c(\mathcal{Z})$  be given and let  $h := g\mathcal{H}^{d-1}|_{\mathcal{Z}} \in \mathcal{M}(\Omega)$ . Then, h is an element of the critical cone  $\mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$ , it holds

(41) 
$$\frac{1}{2} \int_{\mathcal{Z}} g^2 |\nabla \bar{\varphi}| d\mathcal{H}^{d-1} = Q_C^{\bar{x}, \bar{\varphi}}(h),$$

and for every sequence  $\{t_k\} \subset \mathbb{R}^+$  with  $t_k \searrow 0$  there exists a sequence  $\{r_k\} \subset X$  such that  $\bar{x} + t_k h + \frac{1}{2} t_k^2 r_k \in C$  holds for all k and such that  $t_k r_k \stackrel{\star}{\rightharpoonup} 0$ ,  $\|h + \frac{1}{2} t_k r_k\|_X \to \|h\|_X$  and  $\langle \bar{\varphi}, r_k \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} \to Q_{\bar{c}}^{\bar{x}, \bar{\varphi}}(h)$  holds for  $k \to \infty$ .

*Proof.* Since  $\bar{\varphi} \in C_0(\Omega)$  and  $\mathcal{Z} = \{\bar{\varphi} = 0\}$ , it trivially holds  $\langle \bar{\varphi}, h \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} = 0$  and, consequently,  $h \in \bar{\varphi}^{\perp}$ . It remains to show that h is an element of the weak- $\star$  tangent cone  $\mathcal{T}_{\mathcal{C}}^{\star}(\bar{x})$ , that (41) holds, and that for each  $\{t_k\} \subset \mathbb{R}^+$  with  $t_k \searrow 0$  we can find a sequence  $\{r_k\}$  with the desired approximation properties. To prove these three assertions, we proceed in two steps:

Step 1 (Proof in a Rectification Neighborhood): In what follows, we first consider a prototypical situation, where the support of the function g is contained in a rectification neighborhood of the  $C^1$ -manifold  $\mathcal{Z}$ , i.e., in an open set where  $\mathcal{Z}$  resembles

a  $C^1$ -graph. As we will see, in this simplified setting, we can manually construct a sequence  $\{h_k\}$  such that  $r_k := 2(h_k - h)/t_k$  satisfies the conditions in the lemma, see (42), (43), (44) and (45) below.

Let us denote with  $\partial_i$ , i=1,...,d, the partial derivatives of a function and assume that a point  $p \in \mathcal{Z}$ , an open ball  $B \subset \mathbb{R}^{d-1}$ , an open interval J := (a,b), and a map  $\gamma \in C^1(\overline{B})$  are given such that

$$p \in B \times J$$
,  $\overline{B \times J} \subset \Omega$ ,  $\mathcal{Z} \cap (B \times J) = \{(z, \gamma(z)) : z \in B\}$   
 $W := \{(z, z') : z \in B, |z' - \gamma(z)| < \varepsilon\} \subset B \times J$ ,  $\partial_d \bar{\varphi}(p) > 0$ 

for some  $\varepsilon \in (0,1)$ . Let  $0 \le \psi \in C(W)$  and  $g \in C(\mathcal{Z} \cap (B \times J))$  be continuous and bounded functions and let  $t_k \in (0,\infty)$  be a sequence with  $t_k \searrow 0$ . Assume w.l.o.g. that

$$t_k \|g\|_{L^{\infty}} \|\sqrt{1 + |\nabla \gamma|^2}\|_{L^{\infty}} \le \varepsilon$$

for all k (else consider  $\{t_k\}_{k\geq K}$ ,  $K\in\mathbb{N}$  sufficiently large) and extend  $\psi$  by zero outside of W. With some abuse of notation, we extend the sign of g from  $\mathcal{Z}\cap W$  to W by

$$sign(g)(z, z') := sign g(z, \gamma(z)).$$

We further define the sets

$$G_k := \left\{ (z, z') \in B \times J \mid t_k \frac{g^-(z, \gamma(z))}{2} \le \frac{z' - \gamma(z)}{\sqrt{1 + |\nabla \gamma(z)|^2}} \le t_k \frac{g^+(z, \gamma(z))}{2} \right\} \subset W,$$

where  $g^+$  and  $g^-$  are abbreviations for  $\max(0,g)$  and  $\min(0,g)$ , respectively, and set

$$(42) h_k := \frac{2 \operatorname{sign}(g)}{t_k} \mathbb{1}_{G_k},$$

where  $\mathbb{1}_A : \Omega \to \{0,1\}$  denotes the characteristic function of a set  $A \subset \Omega$ . We claim that the above  $h_k$  satisfies

(43) 
$$\lim_{k \to \infty} \left( \int_{\Omega} h_k \psi v d\mathcal{L}^d \right) = \int_{\mathcal{Z} \cap (B \times J)} g \psi v d\mathcal{H}^{d-1} \quad \forall v \in C_0(\Omega),$$

(44) 
$$\lim_{k \to \infty} \left( \int_{\Omega} |h_k| \psi d\mathcal{L}^d \right) = \int_{Z \cap (B \times I)} |g| \psi d\mathcal{H}^{d-1},$$

and

(45) 
$$\lim_{k \to \infty} \left( \int_{\Omega} \frac{2h_k}{t_k} \bar{\varphi} \psi d\mathcal{L}^d \right) = \frac{1}{2} \int_{Z \cap (B \times J)} g^2 \psi |\nabla \bar{\varphi}| d\mathcal{H}^{d-1}.$$

This can be seen as follows: Given a  $v \in C_0(\Omega)$ , we may calculate (using the abbrevi-

ations  $dz := d\mathcal{L}^{d-1}(z)$  and  $dz' := d\mathcal{L}^1(z')$ 

$$\int_{\Omega} h_k \psi v d\mathcal{L}^d = \frac{2}{t_k} \int_{B} \int_{\gamma(z) + t_k}^{\gamma(z) + t_k} \frac{g^+(z, \gamma(z))}{2} \sqrt{1 + |\nabla \gamma(z)|^2} \operatorname{sign}(g) v \psi dz' dz$$

$$= 2 \int_{B} \int_{\frac{g^+(z, \gamma(z))}{2}}^{\frac{g^+(z, \gamma(z))}{2}} \sqrt{1 + |\nabla \gamma(z)|^2} (\operatorname{sign}(g) v \psi)|_{(z, \gamma(z) + t_k z')} dz' dz$$

$$\rightarrow 2 \int_{B} \int_{\frac{g^-(z, \gamma(z))}{2}}^{\frac{g^+(z, \gamma(z))}{2}} \sqrt{1 + |\nabla \gamma(z)|^2} (\operatorname{sign}(g) v \psi)|_{(z, \gamma(z))} dz' dz$$

$$= \int_{B} \sqrt{1 + |\nabla \gamma(z)|^2} (gv \psi)|_{(z, \gamma(z))} dz = \int_{\mathcal{Z} \cap (B \times J)} g \psi v d\mathcal{H}^{d-1}.$$

This yields (43). To obtain (44), we can use exactly the same calculation as above (just replace v with sign(g)). It remains to prove (45). To this end, we compute

$$\begin{split} &\int_{\Omega} \frac{2h_k}{t_k} \bar{\varphi} \psi \mathrm{d}\mathcal{L}^d = \frac{4}{t_k} \int_{B} \int_{\frac{g^{-(z,\gamma(z))}}{2}}^{\frac{g^{+}(z,\gamma(z))}{2}} \sqrt{1+|\nabla\gamma(z)|^2} \left( \mathrm{sign}(g) \bar{\varphi} \psi \right) |_{(z,\gamma(z)+t_k z')} \mathrm{d}z' \mathrm{d}z \\ &= 4 \int_{B} \int_{\frac{g^{-(z,\gamma(z))}}{2}}^{\frac{g^{+}(z,\gamma(z))}{2}} \sqrt{1+|\nabla\gamma(z)|^2} \int_{0}^{1} \partial_d \bar{\varphi}(z,\gamma(z)+st_k z') \mathrm{d}s \left( \mathrm{sign}(g) \psi \right) |_{(z,\gamma(z)+t_k z')} z' \mathrm{d}z' \mathrm{d}z \\ &\to 4 \int_{B} \int_{\frac{g^{-(z,\gamma(z))}}{2}}^{\frac{g^{+}(z,\gamma(z))}{2}} \sqrt{1+|\nabla\gamma(z)|^2} z' \mathrm{d}z' \left( \mathrm{sign}(g) \partial_d \bar{\varphi} \psi \right) |_{(z,\gamma(z))} \mathrm{d}z \\ &= \frac{1}{2} \int_{B} \left( 1+|\nabla\gamma(z)|^2 \right) \left( g^2 \partial_d \bar{\varphi} \psi \right) |_{(z,\gamma(z))} \mathrm{d}z. \end{split}$$

Differentiating  $\bar{\varphi}(z, \gamma(z))$  w.r.t.  $z_i$  yields  $\partial_d \bar{\varphi}(z, \gamma(z)) \partial_i \gamma(z) = -\partial_i \bar{\varphi}(z, \gamma(z))$  for all  $i = 1, \ldots, d-1$ . Thus,

$$\partial_d \bar{\varphi}(z,\gamma(z)) \sqrt{1+|\nabla \gamma(z)|^2} = |\nabla \bar{\varphi}(z,\gamma(z))| \quad \forall z \in B.$$

Hence, we arrive at

$$\begin{split} \int_{\Omega} \frac{2h_k}{t_k} \bar{\varphi} \psi \mathrm{d}\mathcal{L}^d &\to \frac{1}{2} \int_{B} \sqrt{1 + |\nabla \gamma(z)|^2} \, (|\nabla \bar{\varphi}| g^2 \psi)|_{(z, \gamma(z))} \mathrm{d}z \\ &= \frac{1}{2} \int_{\mathcal{Z} \cap (B \times J)} g^2 \psi |\nabla \bar{\varphi}| \mathrm{d}\mathcal{H}^{d-1}. \end{split}$$

This proves that (45) holds and that  $\{h_k\}$  indeed has the desired properties.

Step 2 (Proof in the General Case): In this second part of the proof, we demonstrate that, given an arbitrary but fixed  $h := g\mathcal{H}^{d-1}|_{\mathcal{Z}} \in \mathcal{M}(\Omega), g \in C_c(\mathcal{Z})$ , we can always use a partition of unity and the manifold property of  $\mathcal{Z}$  to reduce the situation to the case studied in Step 1, see (46), (47) and (48) below.

Recall that the implicit function theorem and the definition of  $\mathcal{Z}$  imply that for every  $p \in \mathcal{Z}$  there exist an orthogonal transformation  $R \in O(d)$ , an open ball  $B \subset \mathbb{R}^{d-1}$ , an open interval J := (a, b), and a map  $\gamma \in C^1(\overline{B})$  with values in J such

that

$$\begin{split} p \in R(B \times J), & \overline{R(B \times J)} \subset \Omega, & \mathcal{Z} \cap R(B \times J) = R(\{(z, \gamma(z)) : z \in B\}), \\ W := \{(z, z') : z \in B, |z' - \gamma(z)| < \varepsilon\} \subset B \times J, \\ R(\{(z, z') \in B \times J : \gamma(z) < z'\}) \subset \{\bar{\varphi} > 0\}, \\ R(\{(z, z') \in B \times J : z' < \gamma(z)\}) \subset \{\bar{\varphi} < 0\} \end{split}$$

for some  $\varepsilon > 0$ . Since supp(g) is compact, we may find points  $p_1, \ldots, p_L \in \mathcal{Z}$  with associated  $R_l, B_l$  etc. such that the sets  $R_l(W_l)$ ,  $l = 1, \ldots, L$ , cover supp(g). Define

$$U := \bigcup_{l=1}^{L} R_l(W_l) \subset \Omega$$

and choose a partition of unity  $(\psi_l)_{l=1}^L$  subordinate to the  $R_l(W_l)$ -cover of the set U, i.e., a collection of continuous functions  $\psi_l \in C(U, [0, 1])$  such that

$$\operatorname{supp}(\psi_l) \subset W_l, \quad \sum_l \psi_l = 1 \text{ on } U.$$

Consider now an arbitrary but fixed sequence  $t_k \in (0, \infty)$  with  $t_k \searrow 0$ , extend the functions  $\psi_l$  by zero and define (for k large enough)

$$sign_{l}(g)(R_{l}(z,z')) := sign g(R_{l}(z,\gamma_{l}(z))) \qquad \forall (z,z') \in W_{l}, 
G_{l,k} := \left\{ (z,z') \in B_{l} \times J_{l} \mid \frac{g^{-}(R_{l}(z,\gamma_{l}(z)))}{2} \le \frac{t_{k}^{-1}(z'-\gamma_{l}(z))}{\sqrt{1+|\nabla\gamma_{l}(z)|^{2}}} \le \frac{g^{+}(R_{l}(z,\gamma_{l}(z)))}{2} \right\}, 
h_{k} := \sum_{l=1}^{L} \frac{2 \operatorname{sign}_{l}(g)}{t_{k}} \mathbb{1}_{R_{l}(G_{l,k})} \psi_{l}.$$

Then, it holds  $\bar{x} + t_k h_k \in C = L^{\infty}(\Omega, [0, 1])$  (cf. the signs of the involved functions and  $||h_k||_{L^{\infty}} \leq 2/t_k$ ), and we may deduce from (b) that for all  $v \in C_0(\Omega)$ , we have

(46) 
$$\lim_{k \to \infty} \left( \int_{\Omega} h_k v d\mathcal{L}^d \right) = \sum_{l=1}^L \int_{\mathcal{Z} \cap R_l(B_l \times J_l)} g \psi_l v d\mathcal{H}^{d-1} = \langle v, h \rangle_{C_0(\Omega), \mathcal{M}(\Omega)}$$

(47) 
$$\lim_{k \to \infty} \left( \int_{\Omega} |h_k| d\mathcal{L}^d \right) = \sum_{l=1}^L \int_{\mathcal{Z} \cap R_l(B_l \times J_l)} |g| \psi_l d\mathcal{H}^{d-1} = ||h||_X$$

and

(48) 
$$\lim_{k \to \infty} \left( \int_{\Omega} \frac{2h_k}{t_k} \bar{\varphi} d\mathcal{L}^d \right) = \frac{1}{2} \sum_{l=1}^{L} \int_{\mathcal{Z} \cap R_l(B_l \times J_l)} g^2 \psi_l |\nabla \bar{\varphi}| d\mathcal{H}^{d-1}$$
$$= \frac{1}{2} \int_{\mathcal{Z}} g^2 |\nabla \bar{\varphi}| d\mathcal{H}^{d-1}.$$

The above proves  $h \in \mathcal{T}_C^{\star}(\bar{x})$  and  $h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$ , see (a). From (46), (47) and (48), we obtain further that  $r_k := 2(h_k - h)/t_k$  satisfies  $\bar{x} + t_k h + \frac{1}{2}t_k^2 r_k \in C$  for all k and

$$t_k r_k \stackrel{\star}{\rightharpoonup} 0, \quad \|h + \frac{1}{2} t_k r_k\|_X \to \|h\|_X, \quad \langle \bar{\varphi}, r_k \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} \to \frac{1}{2} \int_{\mathcal{F}} g^2 |\nabla \bar{\varphi}| d\mathcal{H}^{d-1}$$

as  $k \to \infty$ . If we combine this with Proposition 6.9 and Definition 3.1, then the claim follows immediately.

Using Proposition 6.9 and Lemma 6.10, we finally arrive at an explicit formula for the directional curvature functional in the bang-bang case.

THEOREM 6.11. For every tuple  $(\bar{x}, \bar{\varphi}) \in C \times -\mathcal{N}_C^{\star}(\bar{x})$  that satisfies the conditions in Assumption 6.6, it holds

(49) 
$$\begin{cases} h \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi}) \mid Q_C^{\bar{x}, \bar{\varphi}}(h) < \infty \\ = \{ g\mathcal{H}^{d-1}|_{\mathcal{Z}} \mid g \in L^1\left(\mathcal{Z}, \mathcal{H}^{d-1}\right) \cap L^2\left(\mathcal{Z}, |\nabla \bar{\varphi}| \mathcal{H}^{d-1}\right) \}. \end{cases}$$

Moreover, for every element  $h = g\mathcal{H}^{d-1}|_{\mathcal{Z}}$  of the above set, it is true that

(50) 
$$Q_C^{\bar{x},\bar{\varphi}}(h) = \frac{1}{2} \int_{\mathcal{Z}} g^2 |\nabla \bar{\varphi}| d\mathcal{H}^{d-1}.$$

Proof. Proposition 6.9 yields that " $\subset$ " holds in (49) and that " $\geq$ " holds in (50). To obtain the reverse inclusion/inequality, we consider an arbitrary but fixed  $h = g\mathcal{H}^{d-1}|_{\mathcal{Z}} \in \mathcal{M}(\Omega)$  with some  $g \in L^1\left(\mathcal{Z}, \mathcal{H}^{d-1}\right) \cap L^2\left(\mathcal{Z}, |\nabla \bar{\varphi}| \mathcal{H}^{d-1}\right)$ . Using a compact exhaustion of the domain  $\Omega$  and mollification, it is easy to see that we can find a sequence  $(g_n) \subset C_c(\mathcal{Z})$  with  $g_n \to g$  in  $L^1\left(\mathcal{Z}, \mathcal{H}^{d-1}\right) \cap L^2\left(\mathcal{Z}, |\nabla \bar{\varphi}| \mathcal{H}^{d-1}\right)$ . The latter implies that  $h_n := g_n \mathcal{H}^{d-1}|_{\mathcal{Z}}$  satisfies  $h_n \stackrel{\sim}{\rightharpoonup} h$  in X and that

$$\frac{1}{2} \int_{\mathcal{Z}} g_n^2 |\nabla \bar{\varphi}| \mathrm{d}\mathcal{H}^{d-1} \to \frac{1}{2} \int_{\mathcal{Z}} g^2 |\nabla \bar{\varphi}| \mathrm{d}\mathcal{H}^{d-1}.$$

On the other hand, we obtain from Lemma 6.10 that  $h_n \in \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$  holds for all n and that there exist  $\{r_{n,k}\} \subset X$  and  $\{t_{n,k}\} \subset \mathbb{R}^+$  with  $\bar{x} + t_{n,k} h_n + \frac{1}{2} t_{n,k}^2 r_{n,k} \in C$  for all n, k and

$$t_{n,k} \searrow 0, \quad t_{n,k}r_{n,k} \stackrel{\star}{\rightharpoonup} 0, \quad \langle \bar{\varphi}, r_{n,k} \rangle_{C_0(\Omega), \mathcal{M}(\Omega)} \to Q_C^{\bar{x}, \bar{\varphi}}(h_n)$$
  
and  $\|h_n + \frac{1}{2}t_{n,k}r_{n,k}\|_X \to \|h_n\|_X$ 

for all n as  $k \to \infty$ . Note that, since the sequence  $||h_n||_X$  is bounded and since the norms  $||h_n + \frac{1}{2}t_{n,k}r_{n,k}||_X$  converge to  $||h_n||_X$ , in the above situation, we may assume w.l.o.g. that  $||t_{n,k}r_{n,k}||_X \le M$  holds for some constant M that is independent of n and k (just shift the index k appropriately for each n). From Lemmas 3.3 and 6.10, we may now deduce that k is an element of the critical cone  $\mathcal{K}_C^*(\bar{x}, \bar{\varphi})$  and that

$$Q_C^{\bar{x},\bar{\varphi}}(h) \leq \liminf_{n \to \infty} Q_C^{\bar{x},\bar{\varphi}}(h_n) = \liminf_{n \to \infty} \frac{1}{2} \int_{\mathcal{Z}} g_n^2 |\nabla \bar{\varphi}| d\mathcal{H}^{d-1} = \frac{1}{2} \int_{\mathcal{Z}} g^2 |\nabla \bar{\varphi}| d\mathcal{H}^{d-1} < \infty.$$

This proves " $\supset$ " in (49) and " $\leq$ " in (50).

We combine Theorems 6.4 and 6.11 and arrive at the main result of this section.

П

THEOREM 6.12 (Explicit No-Gap Second-Order Condition for Bang-Bang Problems). Consider an optimization problem of the form (P) with C, X etc. as in Assumption 6.2. Assume that  $\bar{x} \in C$  is fixed, that J satisfies the conditions in Assumption 4.1, that the map  $X \ni h \mapsto J''(\bar{x}) h^2 \in \mathbb{R}$  is weak- $\star$  continuous, that  $\bar{x}$  and  $\bar{\varphi}$  satisfy the conditions in Assumption 6.6, and that the constant  $K(\bar{\varphi})$  in (31) is non-zero. Then, the condition

(51) 
$$\frac{1}{2} \int_{\mathcal{Z}} g^{2} |\nabla \bar{\varphi}| d\mathcal{H}^{d-1} + J''(\bar{x}) \left(g\mathcal{H}^{d-1}|_{\mathcal{Z}}\right)^{2} > 0$$
$$\forall g \in L^{1}\left(\mathcal{Z}, \mathcal{H}^{d-1}\right) \cap L^{2}\left(\mathcal{Z}, |\nabla \bar{\varphi}| \mathcal{H}^{d-1}\right) \setminus \{0\}$$

is equivalent to the quadratic growth condition (34) with constants c > 0 and  $\varepsilon > 0$ .

We conclude this section with some remarks.

Remark 6.13.

- (i) The inequality (51) with "\ge " instead of ">" is still a necessary optimality condition when  $K(\bar{\varphi}) = 0$ .
- (ii) Theorem 6.12 is still valid when  $\mathcal{Z} = \emptyset$  and (51) is empty. In this case, the only thing that has to be checked to obtain (34) is the level-set condition  $K(\bar{\varphi}) > 0$ .
- (iii) We expect that the assumptions on  $\bar{\varphi}$  in Theorem 6.12 can be weakened (the Taylor expansion in Lemma 6.7, for example, also holds in a far more general setting, see [12, Proposition 5.9]).
- (iv) The techniques used in the proofs of Lemma 6.8 and Proposition 6.9 might be useable in other situations as well (e.g., the idea to exploit the subdifferential structure of the admissible set C, cf. (40).
- (v) We point out that (MRC) does not hold in the situation of Theorem 6.12 (from  $C \subset L^1(\Omega)$  it follows  $\mathcal{T}_C(\bar{x}) \cap \bar{\varphi}^{\perp} = \{0\} \neq \mathcal{K}_C^{\star}(\bar{x}, \bar{\varphi})$  and this is incompatible with the condition (MRC), cf. Remark 3.5).
- (vi) The first addend on the left-hand side of (51) measures the curvature of the set C in  $\bar{x}$  (compare, e.g., with Theorem 5.11). Note that the surface integral in (51) is only meaningful for a  $C^1$ -function  $\bar{\varphi}$ . This shows that exploiting the regularity of the gradient  $\bar{\varphi} = J'(\bar{x})$  is essential for the derivation of Theorem 6.12.

Finally, we would like to compare our Theorem 6.12 with the results of [11] by means of an example.

Example 6.14. We consider the optimal control problem

(52) 
$$\begin{aligned} & \textit{Minimize} \quad \frac{1}{2} \|y - y_d\|_{L^2(\Omega)}^2 \\ & \textit{such that} \quad -\Delta y + a(y) = u \quad \textit{in } \Omega, \quad y = 0 \quad \textit{on } \partial \Omega \\ & \textit{and} \quad -1 \leq u \leq 1 \quad \textit{in } \Omega. \end{aligned}$$

Here, the state equation is to be understood in the weak sense. We assume that  $\Omega \subset \mathbb{R}^d$ .  $d \in \{1,2,3\}$ , is a bounded domain with Lipschitz boundary  $\partial \Omega$ , that  $y_d \in L^2(\Omega)$  and that  $a: \mathbb{R} \to \mathbb{R}$  is twice continuously differentiable and monotonically increasing. Further, we define the admissible set  $C := L^{\infty}(\Omega; [-1, 1])$ .

Now, we can employ [11, Theorem 2.1] to obtain that the control-to-state operator  $G: L^2(\Omega) \to H^1_0(\Omega) \cap C(\overline{\Omega})$  which maps u to the solution y of the semilinear PDE in (52) is well defined and twice continuously Fréchet differentiable. For  $\bar{u}, h \in L^2(\Omega)$ , the derivative  $z_h := G'(\bar{u})h$  solves the linearized PDE

$$-\Delta z_h + a'(\bar{y})z_h = h \quad \text{in } \Omega, \quad z_h = 0 \quad \text{on } \partial\Omega,$$

with  $\bar{y} = G\bar{u}$ . Consequently, we can check that the reduced objective  $J: L^{\infty}(\Omega) \to \mathbb{R}$ ,  $J(u) = \frac{1}{2} \|Gu - y_d\|_{L^2(\Omega)}^2$  is twice continuously Fréchet differentiable and that the derivatives are given by

$$J'(\bar{u}) h = \int_{\Omega} \bar{\varphi} h \, \mathrm{d}\mathcal{L}^d, \qquad J''(\bar{u}) h^2 = \int_{\Omega} \left[ 1 - a''(\bar{y}) \, \bar{\varphi} \right] z_h^2 \, \mathrm{d}\mathcal{L}^d.$$

Here,  $\bar{\varphi} \in H_0^1(\Omega) \cap C(\bar{\Omega})$  is the solution of the adjoint equation

$$-\Delta \bar{\varphi} + a'(\bar{y})\bar{\varphi} = \bar{y} - y_d \quad \text{in } \Omega, \quad \bar{\varphi} = 0 \quad \text{on } \partial\Omega.$$

In order to apply the previous theorems, we have to check that the Taylor expansion (4) from Assumption 4.1 is satisfied by J with the setting of spaces as in Assumption 6.2. First of all, we mention that the quadratic form  $J''(\bar{u})$  can be extended from  $L^{\infty}(\Omega)^2$  to  $\mathcal{M}(\Omega)^2$  by using results for PDEs with measures on the right-hand side, see the discussion in [11, Section 2.5]. Now, let sequences  $\{h_k\} \subset L^{\infty}(\Omega)$ ,  $\{t_k\} \subset \mathbb{R}^+$  with  $t_k \searrow 0$ ,  $h_k \stackrel{\star}{\longrightarrow} h$  in  $\mathcal{M}(\Omega)$  and  $-1 \le \bar{u} + t_k h_k \le 1$  be given. Then, the twice continuous Fréchet differentiability of  $J: L^{\infty}(\Omega) \to \mathbb{R}$  yields

$$J(\bar{u} + t_k h_k) = J(\bar{u}) + t_k J'(\bar{u}) h_k + \frac{1}{2} t_k^2 J''(\bar{u} + \theta_k t_k h_k) h_k^2$$

for some  $\theta_k \in [0,1]$ . Further, [7, Lemmas 2.6, 2.7] imply that for all  $\varepsilon > 0$ , the inequality

$$\left|J''(\bar{u}) h_k^2 - J''(\bar{u} + \theta_k t_k h_k) h_k^2\right| \le \varepsilon \|z_{h_k}\|_{L^2(\Omega)}^2 \le C \varepsilon \|h_k\|_{L^1(\Omega)}^2 \le \hat{C} \varepsilon$$

holds if k is large enough. Together with the above Taylor expansion, we find

$$\frac{J(\bar{u} + t_k h_k) - J(\bar{u}) - t_k J'(\bar{u}) h_k - \frac{1}{2} t_k^2 J''(\bar{u}) h_k^2}{t_k^2} = \frac{J''(\bar{u} + \theta_k t_k h_k) h_k^2 - J''(\bar{u}) h_k^2}{2}$$

$$\to 0$$

as  $k \to \infty$ . This shows that Assumption 4.1 is satisfied.

In order to proceed, we fix  $\bar{u} \in C$  such that the first-order condition  $J'(\bar{u})(u-\bar{u}) \geq 0$  for all  $u \in C$  is satisfied. We further assume that the adjoint state  $\bar{\varphi}$  has the additional regularity  $\bar{\varphi} \in C^1(\bar{\Omega})$  and we suppose that

$$\min_{x\in N} \lvert \nabla \bar{\varphi}(x) \rvert > 0, \quad \textit{where } N := \{x\in \bar{\Omega} : \bar{\varphi}(x) = 0\}.$$

On the one hand, this condition ensures that Assumption 6.6 holds. On the other hand, from [13, Lemma 3.2] we know that this condition implies

$$\mathcal{L}^d(\{|\bar{\varphi}| \le s\}) \le \hat{K}s \qquad \forall s > 0$$

for some  $\hat{K} > 0$ . From the definition (31), we directly infer  $K(\bar{\varphi}) \geq (4\hat{K})^{-1} > 0$ , such that we can apply Theorems 6.4 and 6.12.

We start by the interpretation of Theorem 6.4 and compare it to the results of [11, 10]. As discussed in Remark 6.5, we obtain that the condition

(54) 
$$J''(\bar{u}) h^2 > -K(\bar{\varphi}) \|h\|_{\mathcal{M}(\Omega)}^2 \quad \forall h \in \mathcal{K}_C^{\star}(\bar{u}, \bar{\varphi}) \setminus \{0\}$$

implies a quadratic growth at  $\bar{u}$  in  $L^1(\Omega)$ , and this sufficient condition is similar to [11, Corollary 2.15]. Here,  $\mathcal{K}_C^{\star}(\bar{u},\bar{\varphi})$  is the critical cone of  $C = L^{\infty}(\Omega;[-1,1])$  w.r.t. the weak-\* topology in  $\mathcal{M}(\Omega)$ , see Definition 2.2. In our situation, we have  $\mathcal{K}_C^{\star}(\bar{u},\bar{\varphi}) = \mathcal{M}(\mathcal{Z})$  with  $\mathcal{Z} = \{x \in \Omega : \bar{\varphi}(x) = 0\}$ , see [11, Corollary 2.13] and (53). Note that [10, Theorem 2.4] improves the constant in [11, Theorem 2.8], and therefore, also [11, Corollary 2.15] can be improved slightly, see also Remark 6.5 (iii). However, it is not possible to obtain a necessary optimality condition similar to (54).

By applying Theorem 6.12, we obtain that

$$(55) \qquad \frac{1}{2} \int_{\mathcal{Z}} g^{2} |\nabla \bar{\varphi}| d\mathcal{H}^{d-1} + J''(\bar{u}) \left(g\mathcal{H}^{d-1}|_{\mathcal{Z}}\right)^{2} > 0 \qquad \forall g \in L^{2}\left(\mathcal{Z}, \mathcal{H}^{d-1}\right) \setminus \{0\}$$

is equivalent to a quadratic growth of J at  $\bar{u}$  w.r.t. the norm in  $L^1(\Omega)$ . Note that due to (53), we do not need to employ weighted Lebesgue spaces on  $\mathcal{Z}$ .

It is interesting to see that the sufficient condition (54) involves all signed finite Radon measures with support contained in  $\mathbb{Z}$ , whereas we only need to consider Radon measures with  $L^2$ -density (w.r.t.  $\mathcal{H}^{d-1}$ ) on  $\mathbb{Z}$  for the no-gap condition (55).

To conclude, our technique of performing a careful analysis of the curvature of the feasible set  $C = L^{\infty}(\Omega; [-1,1]) \subset \mathcal{M}(\Omega)$  allows us to formulate second-order no-gap optimality conditions for the optimal control problem (52). This is not possible with the techniques of [11] which only allow to produce sufficient optimality conditions.

7. Conclusion. Given the examples and applications in Sections 5 and 6, we may conclude that the theoretical framework of Section 4 is indeed a handy tool in the derivation of (no-gap) second-order optimality conditions for problems of the type (P). Our results are not only applicable in those situations where the classical assumptions of polyhedricity and second-order regularity are satisfied (see Theorems 5.10 and 5.11) but also allow to study problems with more complicated admissible sets. This is underlined by the no-gap second-order conditions which were derived in the bangbang setting of Subsection 6.2. Further investigation is needed for the study of the directional curvature functional in the presence of pointwise state constraints, cf. the comments after Example 5.6. We plan to discuss this topic in a forthcoming paper.

## REFERENCES

- [1] L. Ambrosio, Functions of Bounded Variation and Free Discontinuity Problems, Oxford University Press, Oxford & New York, 2000.
- [2] J. Bonnans, Second-order analysis for control constrained optimal control problems of semilinear elliptic systems, Applied Mathematics and Optimization. An International Journal with Applications to Stochastics, 38 (1998), pp. 303-325, https://doi.org/10.1007/ s002459900093.
- [3] J. Bonnans, R. Cominetti, and A. Shapiro, Second order optimality conditions based on parabolic second order tangent sets, SIAM Journal on Optimization, 9 (1999), pp. 466–492, https://doi.org/10.1137/S1052623496306760.
- [4] J. Bonnans and A. Shapiro, Perturbation Analysis of Optimization Problems, Springer Verlag, New York, 2000.
- [5] J. Bonnans and H. Zidani, Optimal control problems with partially polyhedric constraints, SIAM Journal on Control and Optimization, 37 (1999), pp. 1726-1741, https://doi.org/10. 1137/S0363012998333724.
- [6] H. Cartan, Calcul différentiel, Hermann, Paris, 1967.
- [7] E. CASAS, Second order analysis for bang-bang control problems of PDEs, SIAM Journal on Control and Optimization, 50 (2012), pp. 2355-2372, https://doi.org/10.1137/120862892.
- [8] E. CASAS AND F. TRÖLTZSCH, Second order analysis for optimal control problems: improving results expected from abstract theory, SIAM Journal on Optimization, 22 (2012), pp. 261– 279, https://doi.org/10.1137/110840406.
- [9] E. CASAS AND F. TRÖLTZSCH, Second order optimality conditions and their role in PDE control, Jahresbericht der Deutschen Mathematiker-Vereinigung, 117 (2015), pp. 3–44, https://doi. org/10.1365/s13291-014-0109-3.
- [10] E. CASAS, D. WACHSMUTH, AND G. WACHSMUTH, Second order analysis and numerical approximation for bang-bang bilinear control problems, preprint, 2017, https://arxiv.org/abs/1707. 06880.
- [11] E. CASAS, D. WACHSMUTH, AND G. WACHSMUTH, Sufficient second-order conditions for bang-bang control problems, SIAM Journal on Control and Optimization, 55 (2017), pp. 3066–3090, https://doi.org/10.1137/16M1099674.
- [12] C. CHRISTOF AND C. MEYER, Differentiability properties of the solution operator to an elliptic variational inequality of the second kind, Ergebnisberichte des Instituts für Angewandte Mathematik, TU Dortmund, (2016).
- [13] K. Deckelnick and M. Hinze, A note on the approximation of elliptic control problems with bang-bang controls, Computational Optimization and Applications. An International

- Journal, 51 (2012), pp. 931-939, https://doi.org/10.1007/s10589-010-9365-z.
- [14] C. Do, Generalized second-order derivatives of convex functions in reflexive Banach spaces, Transactions of the American Mathematical Society, 334 (1992), pp. 281–301, https://doi. org/10.2307/2153983.
- [15] L. EVANS AND R. GARIEPY, Measure theory and fine properties of functions, CRC Press, Boca Raton, FL, revised ed., 2015.
- [16] P. Grisvard, Elliptic Problems in Nonsmooth Domains, Pitman, Boston, 1985.
- [17] D. HOLM, T. SCHMAH, AND C. STOICA, Geometric Mechanics and Symmetry, Oxford University Press, Oxford, 2009.
- [18] H. MAURER AND J. ZOWE, First and second order necessary and sufficient optimality conditions for infinite-dimensional programming problems, Mathematical Programming, 16 (1979), pp. 98-110, https://doi.org/10.1007/BF01582096.
- [19] P. Mehlitz and G. Wachsmuth, The weak sequential closure of decomposable sets in Lebesgue spaces and its application to variational geometry, Preprint, TU Chemnitz, (2017).
- [20] F. MIGNOT, Controle dans les inéquations variationelles elliptiques, Journal of Functional Analysis, 22 (1976), pp. 130–185.
- [21] R. ROCKAFELLAR AND R. Wets, Variational Analysis, Springer Verlag, Berlin, 1998.
- [22] A. Shapiro, On uniqueness of Lagrange multipliers in optimization problems subject to cone constraints, SIAM Journal on Optimization, 7 (1997), pp. 508-518, https://doi.org/10. 1137/S1052623495279785.
- [23] G. Wachsmuth, A guided tour of polyhedric sets, Preprint, TU Chemnitz, (2018). To appear.
- [24] J. ZOWE AND S. KURCYUSZ, Regularity and stability for the mathematical programming problem in Banach spaces, Applied Mathematics and Optimization, 5 (1979), pp. 49–62, https://doi.org/10.1007/BF01442543.