

**Integration of copper toxicity mechanisms in *Raphidocelis subcapitata*: advancing insights at environmentally relevant concentrations**

Manuela D. Machado<sup>1,2,3</sup> and Eduardo V. Soares<sup>1,2,3\*</sup>

<sup>1</sup>Bioengineering Laboratory, ISEP, Polytechnic of Porto, Rua Dr António Bernardino de Almeida,  
431, 4249-015 Porto, Portugal

<sup>2</sup>CEB - Centre of Biological Engineering, University of Minho, Campus de Gualtar, 4710-057  
Braga, Portugal

<sup>3</sup>LABBELS – Associate Laboratory, Braga/Guimarães, Portugal

**Manuela D. Machado ORCID:** 000-0003-3519-8654


**Eduardo V. Soares ORCID:** 000-0003-2280-5291

**CONTENTS**

Figure S1. Correlation between intracellular ROS accumulation and MDA content in <i>R. subcapitata</i> .....	S2
Figure S2. Correlation between MDA content and % of cells with permeabilized plasma membrane in <i>R. subcapitata</i> exposed to 53 µg/L Cu.....	S2


---

\*Corresponding author email address: [evs@isep.ipp.pt](mailto:evs@isep.ipp.pt)



**Fig. S1** Correlation between intracellular ROS accumulation and MDA content in *R. subcapitata*.

A – cells exposed to 33 µg/L Cu. B – cells exposed to 53 µg/L Cu.



**Fig. S2** Correlation between MDA content and percentage of cells with permeabilized plasma membrane in *R. subcapitata* exposed to 53 µg/L Cu.